

BASES OF T-EQUIVARIANT COHOMOLOGY OF BOTT-SAMELSON VARIETIES

VLADIMIR SHCHIGOLEV

ABSTRACT. We construct combinatorial bases of the T -equivariant cohomology $H_T^\bullet(\Sigma, k)$ of the Bott-Samelson variety Σ under some mild restrictions on the field of coefficients k . This bases allow us to prove the surjectivity of the restrictions $H_T^\bullet(\Sigma, k) \rightarrow H_T^\bullet(\pi^{-1}(x), k)$ and $H_T^\bullet(\Sigma, k) \rightarrow H_T^\bullet(\Sigma \setminus \pi^{-1}(x), k)$, where $\pi : \Sigma \rightarrow G/B$ is the canonical resolution. In fact, we also construct bases of the targets of these restrictions by picking up certain subsets of certain bases of $H_T^\bullet(\Sigma, k)$ and restricting them to $\pi^{-1}(x)$ or $\Sigma \setminus \pi^{-1}(x)$ respectively.

As an application, we calculate the cohomology of the costalk-to-stalk embedding for the direct image $\pi_* k_\Sigma$. This algorithm avoids division by 2, which allows us to reestablish 2-torsion for parity sheaves in Braden's example [BW].

1. INTRODUCTION

Let Σ be a Bott-Samelson variety for a connected semisimple complex group G . In this paper, we study the T -equivariant cohomology $H_T^\bullet(\Sigma, k)$, where T is a maximal torus in G and k is a PID. The direction of our research is mainly determined by Härterich's preprint [H]. However, this preprint uses difficult Arabia's results [A89] and [A98], which, as explicitly stated, are valid for the ring of coefficients $\mathbb{Q}$. Therefore, we prefer not to use geometrical bases (coming from Białynicki-Birula cells) and construct combinatorial bases instead. If the sequence of simple reflections determining Σ has length r , then there are in total 2^{2^r-1} bases B_ρ of $H_T^\bullet(\Sigma, k)$ under some mild restriction on the characteristic of k (Theorem 15 and Lemma 27).

Let $\pi : \Sigma \rightarrow G/B$ be the canonical resolution and $x \in G/B$ be an arbitrary T -fixed point. Using the previously constructed bases of $H_T^\bullet(\Sigma, k)$, we can construct a basis of $H_T^\bullet(\pi^{-1}(x), k)$ as follows (Theorem 17, Remark 19 and Lemma 28):

- (1) choose an index ρ ;
- (2) choose a subset $M \subset B_\rho$;
- (3) consider the restrictions $\{f|_{\pi^{-1}(x)} \mid f \in M\}$.

This fact automatically implies that the restriction $H_T^\bullet(\Sigma, k) \rightarrow H_T^\bullet(\pi^{-1}(x), k)$ is surjective.

One may naturally ask what happens if we consider the compliment $\Sigma \setminus \pi^{-1}(x)$ instead of $\pi^{-1}(x)$? It turns out that there exists a basis $H_T^\bullet(\Sigma \setminus \pi^{-1}(x), k)$ that can be constructed from a basis B_ρ of $H_T^\bullet(\Sigma, k)$ by steps similar to steps (1)–(3) above (Theorem 25, Remark 26 and Lemma 29).

A plausible motivation to consider the T -equivariant cohomology of $\Sigma \setminus \pi^{-1}(x)$ is to calculate the decomposition of the direct image $\pi_* k_\Sigma$ into a direct sum of parity sheaves introduced in [JMW]. It was noted by the authors of this paper that the natural map $i_\lambda^! \mathcal{F} \rightarrow i_\lambda^* \mathcal{F}$ plays a decisive role in determining such a decomposition at least when k is a field (see [JMW, Proposition 2.26]). Here i_λ is the embedding of a (closed) stratum. In this paper, we address the following question:

Problem 1. *Let $\pi : \Sigma \rightarrow G/B$ be a Bott-Samelson resolution and $x \in G/B$ be a T -fixed point. Denote by $i_x : \{x\} \hookrightarrow G/B$ the natural embedding. How to calculate the map $H_T^\bullet(\Sigma, i_x^! \pi_* k_\Sigma) \rightarrow H_T^\bullet(\Sigma, i_x^* \pi_* k_\Sigma)$?*

It is answered in this paper by Corollary 31. Note that unlike [JMW] this problem does not involve any stratifications. However, we can apply its solution to parity sheaves by considering the stratification $G/B = \bigsqcup_{x \in W} BxB/B$ and dividing by the T -equivariant Euler classes of the natural embeddings $\{x\} \hookrightarrow BxB/B$. The corresponding constriction is given in Section 6.6.

Our algorithm is similar to the one described in [S]. It uses the same construction of the transition matrix as in [S, Theorem 4.10.3], which was previously used by Fiebig for his upper bound for Lusztig's conjecture [F] and which originally comes from the same Härterich's preprint [H]. The advantage of our approach here compared with [S] is that we do not divide by 2 when we compute products of the basis elements of $H_T^\bullet(\pi^{-1}(x), k)$ (cf. formula (22) of this paper and [S, Lemma 4.8.3]). This allows us to reestablish the 2-torsion for hexagonal permutations in Braden's example [BW, Appendix A].

The paper is organized as follows. In Section 2, we introduce the main notions: the Bott-Samelson variety, combinatorial galleries, load bearing walls, orders $\triangleleft$ and $<$, tree analogs of combinatorial galleries, etc. In Section 3, we explain how to adjust Brion's proofs [B] of localization theorems to our situation of coefficients different from $\mathbb{C}$ and of noncompact spaces. We use some ideas from [FW], where the authors also prove localization theorems additionally assuming finiteness of T -curves (which is not the case for Bott-Samelson varieties). It is important to notice that we do need some form of GKM-restriction ((C3) in Corollary 6) to prove the intersection formula for the image. To ensure this condition, we first work with coefficients $\mathbb{Z}' = \mathbb{Z}$ or $\mathbb{Z}' = \mathbb{Z}[1/2]$ if the root system contains a component of type C_n and then change coefficients to a PID in Section 6.1.

In Section 4, we construct bases of $H_T^\bullet(\Sigma, \mathbb{Z}')$ and $H_T^\bullet(\Sigma_x, \mathbb{Z}')$. We use here the criterions proved by Härterich [H, Theorems 6.2 and 6.3]. The proofs of these results use only the smooth case of $\mathrm{SL}_2(\mathbb{C})$ or $\mathrm{PSL}_2(\mathbb{C})$, which can be handled by [AB]. We develop here the main combinatorial tool of this paper — operators of copy Δ and concentration ∇_t , which construct elements of $H_T^\bullet(\Sigma, \mathbb{Z}')$ from elements of $H_T^\bullet(\Sigma', \mathbb{Z}')$, where Σ' is the Bott-Samelson variety for the truncated sequence. Finally, we construct bases of $H_T^\bullet(\Sigma_x, \mathbb{Z}')$ in the way similar to [S]. Our new product of bases elements (22) does not include division, which is a definite advantage.

In Section 5, we construct a basis of $H_T^\bullet(\Sigma \setminus \pi^{-1}(x), \mathbb{Z}')$. To achieve this goal, we need to prove the localization theorem for $\Sigma \setminus \pi^{-1}(x)$ and to prove a criterion for $H_T^\bullet(\Sigma \setminus \pi^{-1}(x), \mathbb{Z}')$ (Proposition 21) similar to Härterich's criterions [H, Theorems 6.2 and 6.3].

Section 6 is devoted to applications of the obtained results. We begin with the change of coefficients in Section 6.1, which allows us to obtain bases of $H_T^\bullet(\Sigma, k)$, $H_T^\bullet(\Sigma_x, k)$ and $H_T^\bullet(\Sigma \setminus \pi^{-1}(x), k)$ for any PID k of characteristic not 2 if the root system contains a component of type C_n . Then we solve Problem 1 and in Section 6.6 show how this information can be used to decompose the direct image $\pi_* \underline{k}_\Sigma[r]$ to a direct sum of indecomposable parity sheaves. As an example, we show in Section 6.7 that this decomposition may depend on the characteristic of k (Theorem 37) by considering a hexagonal permutation as in Braden's example [BW, Appendix A].

Finally, we note that all the above results are valid in the affine setting with the corresponding restriction on the characteristic, as we use only local techniques. The reader may consult, for example, [G] about affine pavings.

2. BOTT-SAMELSON VARIETY

Let G be a connected semisimple complex algebraic group, T be its maximal torus and B be its Borel subgroup containing T . We denote by W , Φ , Φ^+ , Π the Weyl group, the set of all roots, the set of positive roots and the set of simple roots respectively.

Let α be a root. We denote by s_α and U_α the simple reflection and the unipotent subgroup corresponding to α respectively. Let G_α be the subgroup of G generated U_α and $U_{-\alpha}$. This subgroup is isomorphic either to $\mathrm{SL}_2(\mathbb{C})$ or to $\mathrm{PSL}_2(\mathbb{C})$. We set $B_\alpha = G_\alpha \cap B$. Let P_α be the parabolic subgroup of G corresponding to α . If α is simple, then $P_\alpha = B \cup Bs_\alpha B$. We denote by $x_\alpha : \mathbb{C} \rightarrow U_\alpha$ the canonical homomorphism.

Throughout the paper, we fix a sequence $s = (s_1, s_2, \dots, s_r)$ of simple reflections, where $s_i = s_{\alpha_i}$ for some $\alpha_i \in \Pi$, and consider the *Bott-Samelson* variety

$$\Sigma = P_{\alpha_1} \times P_{\alpha_2} \times \cdots \times P_{\alpha_r} / B^r,$$

where B^r acts as follows:

$$(p_1, p_2, \dots, p_r) \cdot (b_1, b_2, \dots, b_r) = (p_1 b_1, b_1^{-1} p_2 b_2, \dots, b_{r-1}^{-1} p_r b_r).$$

We denote by $[p_1, \dots, p_r]$ the point of Σ corresponding to $(p_1, \dots, p_r)$. It is well known that Σ is a smooth complex variety of dimension r . Moreover, T acts on Σ by $t \cdot [p_1, p_2, \dots, p_r] = [tp_1, p_2, \dots, p_r]$ and Σ is covered by open T -equivariant subsets

$$U^\gamma = \{[x_{\gamma_1(-\alpha_1)}(c_1)\gamma_1, x_{\gamma_2(-\alpha_2)}(c_2)\gamma_2, \dots, x_{\gamma_r(-\alpha_r)}(c_r)\gamma_r] \mid c_1, c_2, \dots, c_r \in \mathbb{C}\},$$

where γ runs through Γ .

Let $\pi : \Sigma \rightarrow G/B$ be the map $\pi([p_1, \dots, p_r]) = p_1 \cdots p_r B / B$. For any $x \in G/B$, we fix the following notation $\Sigma_x = \pi(x)$ and $\bar{\Sigma}_x = \Sigma \setminus \Sigma_x$. We can also view Σ as a closed subvariety of $(G/B)^r$ via the embedding $\iota : \Sigma \hookrightarrow (G/B)^r$ defined by

$$\iota([p_1, \dots, p_r]) = (p_1 B, p_1 p_2 B, \dots, p_1 p_2 \cdots p_r B).$$

This map is an isomorphism for $G = \mathrm{SL}_2(\mathbb{C})$ and $G = \mathrm{PSL}_2(\mathbb{C})$.

Each point of G/B fixed by T can be written uniquely as wB for some $w \in W$. So, abusing notation, we will denote this point simply by w . Consider the following set:

$$\Gamma = \{(\gamma_1, \dots, \gamma_r) \mid \gamma_i = s_i \text{ or } \gamma_i = e\}.$$

The elements of this set are called *combinatorial galleries*. Note that Γ can be thought of as the set of all T -fixed points of Σ if we identify $(\gamma_1, \dots, \gamma_r)$ with $[\gamma_1, \dots, \gamma_r]$. For each $\gamma = (\gamma_1, \dots, \gamma_r) \in \Gamma$ and $i = 0, \dots, r$, we denote $\gamma^i = \gamma_1 \cdots \gamma_i$. So we get $\gamma^0 = e$. If additionally $i > 0$, then we denote $\beta_i(\gamma) = \gamma^i(-\alpha_i)$ and $\tilde{\beta}_i(\gamma) = \gamma^{i-1}(-\alpha_i)$. If $\beta_i(\gamma) > 0$, then we say that i is *load bearing* for γ or that the wall corresponding to $\beta_i(\gamma)$ is *load bearing*. For any $A \subset W$, we denote

$$\Gamma_A = \{\gamma \in \Gamma \mid \pi(\gamma) \in A\}, \quad \bar{\Gamma}_A = \Gamma \setminus \Gamma_A.$$

If $A = \{x\}$, then we use the simplified notation $\Gamma_x = \Gamma_{\{x\}}$ and $\bar{\Gamma}_x = \bar{\Gamma}_{\{x\}}$.

For $\alpha \in \Phi^+$ and $\gamma \in \Gamma$, we set

$$J(\gamma) = \{i \mid \beta_i(\gamma) > 0\}, \quad M_\alpha(\gamma) = \{i \mid \beta_i(\gamma) = \pm\alpha\}, \quad J_\alpha(\gamma) = \{i \mid \beta_i(\gamma) = \alpha\} = J(\gamma) \cap M_\alpha(\gamma),$$

$$D(\gamma) = \{i \mid \tilde{\beta}_i(\gamma) > 0\}, \quad D_\alpha(\gamma) = \{i \mid \tilde{\beta}_i(\gamma) = \alpha\} = D(\gamma) \cap M_\alpha(\gamma).$$

Note that $\beta_i(\gamma) > 0 \Leftrightarrow \gamma^i s_i < \gamma^i$ and $\tilde{\beta}_i(\gamma) > 0 \Leftrightarrow \gamma^{i-1} s_i < \gamma^{i-1}$. Using these subsets, we introduce the following equivalence relation on Γ :

$$\gamma \sim_\alpha \delta \iff \gamma_i = \delta_i \text{ unless } \beta_i(\gamma) = \pm\alpha.$$

One can easily check that $M_\alpha(\gamma)$ depends only on the $\sim_\alpha$ -equivalence class of γ .

We will use the following two relations on Γ :

$$\begin{aligned} \delta \triangleleft \gamma &\iff \delta^0 = \gamma^0, \dots, \delta^{i-1} = \gamma^{i-1}, \delta^i < \gamma^i \text{ for some } i = 0, \dots, r; \\ \delta < \gamma &\iff \delta^i < \gamma^i, \delta^{i+1} = \gamma^{i+1}, \dots, \delta^r = \gamma^r \text{ for some } i = 0, \dots, r. \end{aligned}$$

Clearly, $\triangleleft$ is a total order on Γ , whereas $<$ in general becomes a total order only when restricted to some Γ_x . As usual, we set $\delta \trianglelefteq \gamma \Leftrightarrow \delta \triangleleft \gamma$ or $\delta = \gamma$ and $\delta \leq \gamma \Leftrightarrow \delta < \gamma$ or $\delta = \gamma$. Note that $\delta \sim_\alpha \gamma$ and $J_\alpha(\delta) \subset J_\alpha(\gamma)$ imply $\delta \trianglelefteq \gamma$. We recall the following Lemma from [H].

Proposition 1. *Let $M_\alpha(\gamma) = \{i_1 < \dots < i_\ell\}$. Then for $1 \leq j < \ell$ we have*

$$i_j \in J_\alpha(\gamma) \Leftrightarrow i_{j+1} \in D_\alpha(\gamma).$$


and $i_\ell \in J_\alpha(\gamma)$ if and only if $s_\alpha \pi(\gamma) < \pi(\gamma)$. In particular, if $\gamma \sim_\alpha \delta$ and $\pi(\gamma) = \pi(\delta)$, then $J_\alpha(\delta) \subset J_\alpha(\gamma) \Leftrightarrow D_\alpha(\delta) \subset D_\alpha(\gamma)$.

We use the symbol $\cdot$ to denote the addition of a new entry to a sequence: $(a_1, \dots, a_n) \cdot b = (a_1, \dots, a_n, b)$. Conversely, for a nonempty sequence $a = (a_1, \dots, a_n)$, we denote $a' = (a_1, \dots, a_{n-1})$ its truncation.

For any integer $r \geq 0$, let Tr_r denote the binary tree that consisting of all sequences (including the empty one) with entries 0 or 1 of length less than r . We obviously have $|\text{Tr}_r| = 2^r - 1$ and $\text{Tr}_r = \text{Tr}_{r-1} \cdot 0 \sqcup \text{Tr}_{r-1} \cdot 1 \sqcup \{\emptyset\}$ for $r > 0$. To construct bases of the T -equivariant cohomology of the Bott-Samelson varieties, we consider the following set:

$$\Upsilon = \{\rho : \text{Tr}_r \rightarrow \{s_1, \dots, s_r\} \mid \rho_u = e \text{ or } \rho_u = s_{r-|u|}\}.$$

Elements of this set are thus tree analogs of combinatorial galleries. For example, for $r = 3$, we draw Υ as follows:



where the right column shows the sets to which the elements of the corresponding rows belong.

Our notation above implicitly referred to the sequence $s = (s_1, \dots, s_r)$ and the group G . If, for the sake of induction, we want to consider the same objects for the shorter sequence $s' = (s_1, \dots, s_{r-1})$, we add $'$ to our symbols: Σ' , Γ' , $\bar{\Gamma}'_x$, etc. For example, let $r > 0$ and $\rho \in \Upsilon$. For any $\varepsilon \in \{0, 1\}$, we consider its ε -truncation $\rho'_\varepsilon \in \Upsilon'$ defined by $(\rho'_\varepsilon)_{u'} = \rho_{u' \cdot \varepsilon}$ (at the picture above, ρ'_0 and ρ'_1 are the left and right subtrees respectively).

In Section 5, we consider the cases $G = \text{SL}_2(\mathbb{C})$ and $G = \text{PSL}_2(\mathbb{C})$. The sequence s is characterized then only by its length r . We denote by Σ_r^2 and Γ_r^2 the Bott-Samelson variety and the set of combinatorial galleries respectively.

Finally, note that we also consider the Bott-Samelson variety corresponding to the empty sequence ($r = 0$). This is the one-point variety $\Sigma = \Gamma = \{\emptyset\}$.

3. LOCALIZATION THEOREMS

3.1. Set theoretical notation. We denote the fact that N is a subset of M , including the case $N = M$, by $N \subset M$, reserving the notation $N \subsetneq M$ for the proper inclusion. We write $i_{M,N} : N \hookrightarrow M$ for the natural inclusion map. We sometimes write $r_{N,M}^\bullet$ for the map $H_G^\bullet(M, k) \rightarrow H_G^\bullet(N, k)$ induced by a G -equivariant embedding $i_{M,N} : N \hookrightarrow M$. We denote by $|X|$ the cardinality of a finite set X and by $\text{Map}(X, Y)$ the set of all maps from X to Y . For a set S with an equivalence relation $\sim$, we denote by $\text{rep}(S, \sim)$ any set of representatives of $\sim$ -equivalence classes. Throughout the paper, “graded” means “ $\mathbb{Z}$ -graded”.

3.2. Isomorphism of localizations of modules. We want to formulate here a simple lemma from commutative algebra whose prove is left to the reader.

Let S be a (unitary) commutative ring, M and N be S -modules and $q \in S$. Consider the ring of quotients $S' = S[q^{-1}]$ and S' -modules of quotients $M' = M[q^{-1}]$ and $N' = N[q^{-1}]$. Any homomorphism of S -modules $f : M \rightarrow N$ gives rise to the homomorphism $f' : M' \rightarrow N'$ of S' -modules acting as follows: $f'(m/q^k) = f(m)/q^k$.

Lemma 2. *Suppose that for some integers $a, b \geq 0$ the following conditions hold:*

- (1) $q^a N \subset \text{im } f$;
- (2) $q^b \ker f = 0$.

Then $f' : M' \rightarrow N'$ is an isomorphism of S' -modules.

3.3. The equivariant Mayer-Vietoris sequence for open subsets. In this paper, we consider the equivariant derived category $D_T^b(X, k)$ for a commutative ring k and a topological group T . For any object $\mathcal{F}$ of this category, one can define the T -equivariant cohomology $H_T^\bullet(X, \mathcal{F})$. The basic definitions and properties of this category and T -equivariant cohomologies can be found in [BL]. We shall also use the functors f_* , f^* , $f_!$, $f^!$ between equivariant derived categories defined in this book. Remember the following well-known result.

Proposition 3 (Mayer-Vietoris sequence). *Let X be a T -space. For any open T -stable subsets U, V and an object $\mathcal{F} \in D_T^b(X, k)$, we have the following exact sequence:*

$$\begin{aligned} \cdots \rightarrow H_T^{i-1}(X, \mathcal{F}|_{U \cap V}) \rightarrow H_T^i(X, \mathcal{F}|_{U \cup V}) \rightarrow H_T^i(U, \mathcal{F}|_U) \oplus H_T^i(V, \mathcal{F}|_V) \rightarrow \\ \rightarrow H_T^i(X, \mathcal{F}|_{U \cap V}) \rightarrow H_T^{i+1}(X, \mathcal{F}|_{U \cup V}) \rightarrow \cdots \end{aligned}$$

In the proofs of the localization theorems, this proposition is applied as follows. Suppose that $X = U \cup V$, where X is a T -space and U, V are its open T -stable subspaces. Suppose additionally that there exist elements $u \in H_T^n(\text{pt}, k)$ and $v \in H_T^m(\text{pt}, k)$ such that u annihilates $H_T^\bullet(O_U, \mathcal{F}_U)$ and v annihilates $H_T^\bullet(O_V, \mathcal{F}_V)$ for any open T -stable subsets $O_U \subset U$, $O_V \subset V$ and any objects $\mathcal{F}_U \in D_T^b(O_U, k)$, $\mathcal{F}_V \in D_T^b(O_V, k)$. Then Proposition 3 implies that u^2v and uv^2 annihilate $H_T^\bullet(O, \mathcal{F})$ for any open $O \subset X$ and object $\mathcal{F} \in D_T^b(O, k)$.

Another trivial corollary of the Mayer-Vietoris sequence is as follows.

Corollary 4. *Let X be a T -space, $X = \bigsqcup_{i \in I} X_i$, each X_i be open and T -stable and I be finite. Suppose that $Y \subset X$ is another T -subspace. We denote $Y_i = Y \cap X_i$. An element $f \in H^\bullet(Y, k)$ belongs to the image of the restriction $H_T^\bullet(X, k) \rightarrow H_T^\bullet(Y, k)$ if and only if each $f|_{Y_i}$ belongs to image of the restriction $H_T^\bullet(X_i, k) \rightarrow H_T^\bullet(Y_i, k)$.*

Proof. Induction together with the finiteness of I reduces the problem to the case $I = \{1, 2\}$. As $X_1 \cap X_2 = \emptyset$ and the Mayer-Vietoris sequence is compatible with restrictions, we get the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_T^\bullet(X, k) & \longrightarrow & H_T^\bullet(X_1, k) \oplus H_T^\bullet(X_2, k) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & H_T^\bullet(Y, k) & \longrightarrow & H_T^\bullet(Y_1, k) \oplus H_T^\bullet(Y_2, k) & \longrightarrow & 0 \end{array}$$

Hence the required result follows. □

The above arguments apply to any topological group T not necessarily a torus. In this paper, we are however interested only in the case of a torus and use the following notation:

$$S_k = H_T^\bullet(\text{pt}, k) \simeq S(X(T) \otimes_{\mathbb{Z}} k),$$

where $X(T)$ is the character group of T and S in the right-hand side means taking the symmetric algebra. This is a graded algebra such that $S_k^2 = X(T) \otimes_{\mathbb{Z}} k$. Finally, note that in the next section, we need to consider the compact subtorus $K = (S^1)^n$ of $T \simeq \mathbb{C}^n$. We can replace T -equivariant cohomology with K -equivariant cohomology if necessary.

3.4. Localization. We prove here some localization theorems, closely following [B] (see also [FW] for the case of coefficients different from $\mathbb{C}$).

Theorem 5. *Let $\Gamma < T$ be a closed subgroup of T and X be a paracompact T -space that has an open covering $X = \bigcup_{i \in I} Y^{(i)}$ such that for any $i \in I$*

- $Y^{(i)}$ is open and T -equivariant;
- there exists a T -equivariant embedding of $Y^{(i)}$ in a finite dimensional rational representation $V^{(i)}$ of T .

Denote Λ_Γ the set of all weights of T occurring as weights of some $V^{(i)}$ and having nontrivial restriction to Γ .

Then the natural restriction morphism $H_T^\bullet(X, k) \rightarrow H_T^\bullet(X^\Gamma, k)$ becomes an isomorphism after inverting all elements of $\Lambda_\Gamma \otimes_{\mathbb{Z}} k$.

Proof. For simplicity of notation, we assume that $Y^{(i)}$ is a subset of $V^{(i)}$. Let us write

$$V^{(i)} = \mathbb{C}_{\lambda_1^{(i)}} \oplus \cdots \oplus \mathbb{C}_{\lambda_{n_i}^{(i)}},$$

where $\mathbb{C}_\lambda$ is the representation of T with weight $\lambda \in X(T)$. Let U be an open K -invariant neighbourhood of X^Γ in X . Then the set $Y^{(i)} \setminus U$ does not have Γ -fixed points, which we prefer to write as

$$(Y_i \setminus U) \cap (V^{(i)})^\Gamma = \emptyset. \quad (1)$$

Without loss of generality, we can assume that $\lambda_j^{(i)}$ restricts trivially to Γ if and only if $j \leq m_i$. Then $(V^{(i)})^\Gamma$ consists of the points $(c_1, \dots, c_{m_i}, 0, \dots, 0)$. Consider the open subsets $W_j^{(i)} = \{(c_1, \dots, c_{n_i}) \in V^{(i)} \mid c_j \neq 0\}$ of $V^{(i)}$. It follows from (1) that

$$Y_i \setminus U \subset \bigcup_{j=m_i+1}^{n_i} W_j^{(i)}.$$

For any set of the union in the right-hand side, there exists a T -equivariant map $W_j^{(i)} \rightarrow \mathbb{C}_{\lambda_j^{(i)}}^\times \cong T / \ker \lambda_j^{(i)}$, which is the projection to the j^{th} coordinate.

Let us take an open K -invariant subset $O \subset (Y^{(i)} \setminus U) \cap W_j^{(i)}$ for $j > m_i$. Then composition of maps $O \hookrightarrow W_j^{(i)} \rightarrow \mathbb{C}_{\lambda_j^{(i)}}^\times \rightarrow \text{pt}$ gives rise to the following sequence of cohomologies:

$$H_K^2(\text{pt}, k) \rightarrow H_K^2(\mathbb{C}_{\lambda_j^{(i)}}^\times, k) \rightarrow H_K^2(W_j^{(i)}, k) \rightarrow H_K^2(O, k).$$

Identifying T -equivariant and K -equivariant cohomologies, we obtain that the image of the first Chern class $c_1(\lambda_j^{(i)}) \otimes k$ is zero (already for the first map as follows from [J, §1.9(1)]). Writing this Chern class as $\lambda_j^{(i)} \otimes k$, we get that it annihilates $H_K^\bullet(O, \mathcal{F})$ for any $\mathcal{F} \in D_K^b(O, k)$.

Gluing all subsets $(Y^{(i)} \setminus U) \cap W_j^{(i)}$ by the Mayer-Vietoris sequence for open subsets by the method described in Section 3.3, we get the following property:

$$\text{there exist naturals } a_i \text{ such that } q = \prod_{i \in I} \prod_{j=m_i+1}^{n_i} (\lambda_j^{(i)} \otimes k)^{a_i} \text{ annihilates } H_K^\bullet(X \setminus U, \mathcal{F}) \quad (2)$$

for any object $\mathcal{F} \in D_K^b(X \setminus U, k)$.

Consider the following direct limit $L^n(k) := \lim_{\rightarrow U \supset X^\Gamma} H_K^n(U, k)$ that runs over all K -invariant open neighbourhoods U of X^Γ . Denote by $\alpha_U^n : H_K^n(U, k) \rightarrow L^n(k)$ its natural morphisms. We define $L^\bullet(k) = \bigoplus_{n \in \mathbb{Z}} L^n(k)$. It is an S_k -module and we get homomorphisms $\alpha_U^\bullet : H_K^\bullet(U, k) \rightarrow L^\bullet(k)$ of S_k -modules.

We are going to apply Lemma 2 to prove that $\alpha_X^\bullet$ becomes an isomorphism after inverting q . We know that $q \in S_k^{2t}$ for some $t \in \mathbb{Z}$.

Let us check condition (1) of Lemma 2. Let $u \in L^n(k)$. By the definition of the direct limit $u = \alpha_U^n(\bar{u})$ for some $\bar{u} \in H_K^n(U, k)$ and some K -invariant open U containing X^Γ . We have the following exact sequence

$$H_K^\bullet(X, k) \xrightarrow{r_{\bar{U}, X}^\bullet} H_K^\bullet(U, k) \xrightarrow{\partial^\bullet} H_K^{\bullet+1}(X \setminus U, i^! \underline{k}_X),$$

where $i : X \setminus U \hookrightarrow X$ is the natural embedding. Hence and from (2), we get $\partial^{n+2t}(q\bar{u}) = q\partial^n(\bar{u}) = 0$. The exactness of the above sequence yields $q\bar{u} = r_{U, X}^{n+2t}(v)$ for some $v \in H_K^{n+2t}(X, k)$. It remains to recall the following commutative diagram:

$$\begin{array}{ccc} & & L^\bullet(k) \\ & \nearrow \alpha_X^\bullet & \uparrow \alpha_U^\bullet \\ H_K^\bullet(X, k) & & H_K^\bullet(U, k) \\ & \searrow r_{\bar{U}, X}^\bullet & \end{array}$$

from the definition of the direct limit and write

$$qu = q\alpha_U^n(\bar{u}) = \alpha_U^{n+2t}(q\bar{u}) = \alpha_U^{n+2t} \circ r_{U, X}^{n+2t}(v) = \alpha_X^{n+2t}(v).$$

Let us check now condition (2) of Lemma 2. Take some $v \in H_K^n(X, k)$ such that $\alpha_X^n(v) = 0$. By the definition of the direct limit, we get $v|_U = 0$ for some K -invariant open U containing X^Γ . The exactness of the sequence

$$H_K^\bullet(X \setminus U, i^! \underline{k}_X) \xrightarrow{\beta_U^\bullet} H_K^\bullet(X, k) \xrightarrow{r_{\bar{U}, X}^\bullet} H_K^\bullet(U, k)$$

implies $v = \beta_U^n(w)$ for some $w \in H_K^\bullet(X \setminus U, i^! \underline{k}_X)$. Multiplying by q and applying (2), we get $qv = \beta_U^{n+2t}(qw) = 0$.

The universal mapping property for direct limits yields the (unique) morphism $\gamma^\bullet$ such that the diagram

$$\begin{array}{ccc} L^\bullet(k) & \xleftarrow{\quad \gamma^\bullet \quad} & H_K^\bullet(X^\Gamma, k) \\ & \nwarrow \alpha_U^\bullet & \nearrow r_{X^\Gamma, U}^\bullet \\ & H_K^\bullet(U, k) & \end{array}$$

is commutative for any open U containing X^Γ . By [Q, (1.9)], $\gamma^\bullet$ is an isomorphism. It is obviously an isomorphism of S_k -modules. Considering the case $U = X$ and applying the fact that $\alpha_X^\bullet$ becomes an isomorphism after inverting q , we get that $r_{X^\Gamma, X}^\bullet$ also becomes an isomorphism after inverting q and moreover after inverting all elements of $\Lambda_\Gamma \otimes_{\mathbb{Z}} k$. $\square$

As our next step, we explain how to adjust Theorem 6 from Brion's paper [B] to the case of arbitrary coefficients.

Corollary 6 (cf. [B, Theorem 6]). *Let $H_T^\bullet(X, k)$ be a free S_k -module. Under the hypothesis of Theorem 5 with the additional assumption that $\Lambda_T \otimes \mathbf{1}_k$ does not contain zero divisors of S_k , the restriction*

$$i_{X, X^T}^* : H_T^\bullet(X, k) \rightarrow H_T^\bullet(X^T, k)$$

is an embedding.

Moreover, if $H_T^\bullet(X^T, k)$ does not have S_k -torsion (for example, X^T is finite) and the following conditions hold:

- (C1) *k is a unique factorization domain;*
- (C2) *$\lambda \otimes \mathbf{1}_k$ is prime in $S(X(T) \otimes_{\mathbb{Z}} k)$ for any $\lambda \in \Lambda_T$;*
- (C3) *$\lambda \otimes \mathbf{1}_k \notin \Lambda_{\ker \lambda} \otimes_{\mathbb{Z}} k$ for any $\lambda \in \Lambda_T$,*

then we have

$$\operatorname{im} i_{X, X^T}^* = \bigcap_{\lambda \in \Lambda_T} \operatorname{im} i_{X^{\ker \lambda}, X^T}^*.$$

Proof. Let $M = H_T^\bullet(X, k)$, $N = H_T^\bullet(X^T, k)$, $R = \Lambda_T \otimes_{\mathbb{Z}} k$, $S = S_k$, $S' = R^{-1}S$, $M' = R^{-1}M$, $N' = R^{-1}N$, $\varphi = i_{X, X^T}^*$ and φ' be the morphism from M' to N' induced by φ . By Theorem 5, φ' is an isomorphism.

Let $\{e_j\}_{j \in J}$ be an S -basis of M . Then $\{e_j/1\}_{j \in J}$ is an S' -basis of M' . Suppose that $\varphi(\alpha_1 e_{j_1} + \dots + \alpha_k e_{j_k}) = 0$ for $\alpha_1, \dots, \alpha_k \in S$ and mutually distinct indices $j_1, \dots, j_k \in J$. We get

$$\varphi' \left(\frac{\alpha_1}{1} \cdot \frac{e_{j_1}}{1} + \dots + \frac{\alpha_k}{1} \cdot \frac{e_{j_k}}{1} \right) = \varphi' \left(\frac{\alpha_1 e_{j_1} + \dots + \alpha_k e_{j_k}}{1} \right) = \frac{\varphi(\alpha_1 e_{j_1} + \dots + \alpha_k e_{j_k})}{1} = 0.$$

Hence $\alpha_1/1 = \dots = \alpha_k/1 = 0$ in S' . Therefore $\alpha_1 = \dots = \alpha_k = 0$, as R does not contain zero divisors.

Now let us prove the second statement. Let $e_j^* : M \rightarrow S$ and $(e')_j^* : M' \rightarrow S'$ be the j^{th} coordinate functions for M and M' , respectively. Consider the commutative diagram

$$\begin{array}{ccccc} S & \xleftarrow{e_j^*} & M & \xrightarrow{\varphi} & N \\ \downarrow \iota & & \downarrow & \nearrow & \downarrow \\ S' & \xleftarrow{(e')_j^*} & M' & \xleftarrow{(\varphi')^{-1}} & N' \end{array}$$

Denoting the dashed arrow by f_j , we get the following relation:

$$f_j \circ \varphi = \iota \circ e_j^*. \quad (3)$$

Note that all functions f_j uniquely define elements of N :

$$f_j(u) = f_j(u') \quad \forall j \in J \implies u = u'. \quad (4)$$

Let us take $u \in \bigcap_{\lambda \in \Lambda_T} \operatorname{im} i_{X^{\ker \lambda}, X^T}^*$. Consider the coefficients $f_j(u) \in S'$. If they all belong to $\iota(S)$, then in view of (3), the following calculation is possible:

$$f_j \circ \varphi \left(\sum_{j \in J} \iota^{-1}(f_j(u)) e_j \right) = \iota \circ e_j^* \left(\sum_{j \in J} \iota^{-1}(f_j(u)) e_j \right) = f_j(u).$$

Now (4) implies that $u = \varphi \left(\sum_{j \in J} f_j(u) e_j \right) \in \operatorname{im} i_{X, X^T}^*$.

It only remains to prove that $f_j(u) \in S$ for all $j \in J$. Suppose the contrary holds. By (C2), in this case $f_j(u)$ contains an uncancellable prime denominator $\lambda \otimes \mathbf{1}_k$ for some $j \in J$ and $\lambda \in \Lambda_T$.

To proceed, let us introduce the following notation: $\Gamma = \ker \lambda$, $N_\lambda = H_T^\bullet(X^\Gamma, k)$, $R_\lambda = \Lambda_\Gamma \otimes_{\mathbb{Z}} k$, $S'_\lambda = R_\lambda^{-1}S$, $M'_\lambda = R_\lambda^{-1}M$, $N'_\lambda = R_\lambda^{-1}N_\lambda$, $\varphi_\lambda = i_{X, X^\Gamma}^*$ and φ'_λ be the morphism from M'_λ to N'_λ induced by φ_λ . By theorem 5, φ'_λ is an isomorphism.

As $u \in \text{im } i_{X^\Gamma, X^T}^*$, we can write $u = i_{X^\Gamma, X^T}^*(v)$ for some $v \in N_\lambda$. Similarly to the diagram above, we have the following commutative diagram:

$$\begin{array}{ccc} M & \xrightarrow{\varphi_\lambda} & N_\lambda \\ \downarrow & & \downarrow \\ M'_\lambda & \xleftarrow[\varphi'_\lambda]{\sim} & N'_\lambda \end{array}$$

There exists some product $\mathcal{P}_\lambda$ of elements of R_λ such that $(\mathcal{P}_\lambda/1)(\varphi'_\lambda)^{-1}(v/1) = m/1$ for some $m \in M$. Applying φ'_λ to this equality, we get $\mathcal{P}_\lambda v/1 = (\varphi'_\lambda)(m/1) = \varphi_\lambda(m)/1$, which is an equality in N'_λ . Therefore, there exists another product $\mathcal{P}'_\lambda$ of elements of R_λ such that

$$\mathcal{P}'_\lambda \mathcal{P}_\lambda v = \mathcal{P}'_\lambda \varphi_\lambda(m) = \varphi_\lambda(\mathcal{P}'_\lambda m).$$

Applying i_{X^Γ, X^T}^* to the both sides of this equality, we get

$$\mathcal{P}'_\lambda \mathcal{P}_\lambda u = i_{X^\Gamma, X^T}^*(\mathcal{P}'_\lambda \mathcal{P}_\lambda v) = i_{X^\Gamma, X^T}^* \circ \varphi_\lambda(\mathcal{P}'_\lambda m) = \varphi(\mathcal{P}'_\lambda m).$$

Finally applying f_j , we get

$$(\mathcal{P}'_\lambda \mathcal{P}_\lambda/1)f_j(u) = f_j(\mathcal{P}'_\lambda \mathcal{P}_\lambda u) = f_j \circ \varphi(\mathcal{P}'_\lambda m) = e_j^*(\mathcal{P}'_\lambda m)/1 \in \iota(S).$$

This is a contradiction, as $\mathcal{P}'_\lambda \mathcal{P}_\lambda$ by our GKM-restriction (C3) does not have factors proportional to $\lambda \otimes \mathbf{1}_k$. $\square$

3.5. Case of Bott-Samelson varieties. In what follows, we shall only consider the case where the ring of coefficients k is a PID of characteristic not equal to 2 if the root system contains a component of type C_n . As the ordinary cohomology $H^\bullet(\Sigma, k)$ vanishes in odd degrees and is a free k -module in each degree, the degeneracy of the Leray spectral sequence at the E_2 -term implies

$$H_T^\bullet(\Sigma, k) \simeq H^\bullet(\Sigma, k) \otimes_k S_k.$$

Therefore, we can apply the first part of Corollary 6 to prove that the restriction morphism $H_T^\bullet(\Sigma, k) \rightarrow H_T^\bullet(\Gamma, k)$ is an embedding. We denote its image by $\mathcal{X}(k)$.

Similarly, $H_T^\bullet(\Sigma_x, k) \simeq H^\bullet(\Sigma_x, k) \otimes_k S_k$ and we can apply Corollary 6 to prove that the restriction morphism $H_T^\bullet(\Sigma_x, k) \rightarrow H_T^\bullet(\Gamma_x, k)$ is an embedding. We denote its image by $\mathcal{X}^x(k)$.

In order to ensure conditions (C1)–(C3) of Corollary 6, we want to fix ring $\mathbb{Z}'$ for each root system as follows: $\mathbb{Z}' = \mathbb{Z}[1/2]$ if the root system contains a component of type C_n and $\mathbb{Z}' = \mathbb{Z}$ otherwise. This choice automatically guarantees that Theorem 5 and Corollary 6 hold for $k = \mathbb{Z}'$, since $\Lambda_T \subset \Phi$ in these assertions.

Therefore, from now on, we will assume that the cohomologies (ordinary and equivariant) are taken with coefficients $\mathbb{Z}'$ unless otherwise explicitly stated. We also set $S = S_{\mathbb{Z}'}$, $\mathcal{X} = \mathcal{X}(\mathbb{Z}')$ and $\mathcal{X}^x = \mathcal{X}^x(\mathbb{Z}')$.

4. BASES OF THE IMAGES $\mathcal{X}$ AND $\mathcal{X}^x$

4.1. Härterich's localization theorems. We formulate here the following two results due to Härterich [H]. It is important to note that one needs to be more careful with the ring of coefficients when applying the localization theorems in the proofs of these results. For our ring of coefficients $\mathbb{Z}'$, one can apply Theorem 5 and Corollary 6.

Proposition 7 ([H, Theorem 6.2]). *An element $f \in H_T^\bullet(\Gamma)$ belongs to the image $\mathcal{X}$ of the restriction $i_{\Sigma, \Gamma}^* : H_T^\bullet(\Sigma) \rightarrow H_T^\bullet(\Gamma)$ if and only if*

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}$$

for any positive root α and gallery $\gamma \in \Gamma$.

Proposition 8 ([H, Theorem 6.3]). *An element $f \in H_T^\bullet(\Gamma_x)$ belongs to the image $\mathcal{X}^x$ of the restriction $i_{\Sigma_x, \Gamma_x}^* : H_T^\bullet(\Sigma_x) \rightarrow H_T^\bullet(\Gamma_x)$ if and only if*

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, D_\alpha(\delta) \subset D_\alpha(\gamma)} (-1)^{|D_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}}$$

for any positive root α and gallery $\gamma \in \Gamma_x$.

The reader can find the proofs of these results either in Härterich's original preprint [H] or derive them from the proof of Proposition 21 by similarity.

4.2. Copy and concentration. In this section, we describe two ways to get elements of $\mathcal{X}$ from elements of $\mathcal{X}'$. Suppose that $r > 0$. For $f' \in H_T^\bullet(\Gamma')$, we define its *copy* $\Delta f' \in H_T^\bullet(\Gamma)$ by $\Delta f'(\gamma) = f'(\gamma')$ for any $\gamma \in \Gamma$. Clearly, Δ is an S -linear operation.

Lemma 9. $\Delta f' \in \mathcal{X}$ if $f' \in \mathcal{X}'$.

Proof. By Proposition 7, we must prove that

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}} \quad (5)$$

for any $\gamma \in \Gamma$ and $\alpha \in \Phi^+$. We shall use the notation $M_\alpha(\gamma) = \{i_1 < \dots < i_\ell\}$ and $x = \pi(\gamma)$.

Case 1: $r \notin M_\alpha(\gamma)$. In this case, $\delta \sim_\alpha \gamma$ implies $\delta_r = \gamma_r$. Therefore, we can rewrite (5) as follows

$$\sum_{\delta' \in \Gamma', \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|}}, \quad (6)$$

which holds by Proposition 7 applied to $f' \in \mathcal{X}'$.

Case 2: $r \in M_\alpha(\gamma) \setminus J_\alpha(\gamma)$. Choosing in (5) the gallery δ so that $r \notin J_\alpha(\delta)$, we can rewrite this equivalence as (6).

Case 3: $r \in J_\alpha(\gamma)$. Consider the following equivalence relation on the set $\{\delta \in \Gamma \mid \delta \sim_\alpha \gamma\}$: $\delta \equiv \tau \Leftrightarrow \delta' = \tau'$. Clearly, every equivalence class of this relation consists of exactly two elements. Therefore the sum in (5) can be broken into a sum of the following subsums:

$$(-1)^{|J_\alpha(\delta)|} f'(\delta') + (-1)^{|J_\alpha(\tau)|} f'(\tau')$$

for different $\delta \equiv \tau$. As $|J_\alpha(\delta)|$ and $|J_\alpha(\tau)|$ have different parities, the above sum equals zero. $\square$

For $f' \in H_T^\bullet(\Gamma')$ and $t \in \{e, s_r\}$, we define $\nabla_t f' \in H_T^\bullet(\Gamma)$ called the *concentration* of f' at t by

$$\nabla_t f'(\gamma) = \begin{cases} \beta_r(\gamma) f'(\gamma') & \text{if } \gamma_r = t; \\ 0 & \text{otherwise.} \end{cases}$$

for any $\gamma \in \Gamma$. Clearly, ∇_t is an S -linear operation.

Lemma 10. $\nabla_t f' \in \mathcal{X}$ if $f' \in \mathcal{X}'$.

Proof. We shall write down the proof for $\nabla_e f'$, the proof for $\nabla_{s_r} f'$ being similar.

By Proposition 7, we must prove that

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, \delta_r = e, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} \beta_r(\delta) f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}} \quad (7)$$

for any $\gamma \in \Gamma$ and $\alpha \in \Phi^+$. Clearly, it suffices to consider the case $J_\alpha(\gamma) \neq \emptyset$. We shall use the notation $M_\alpha(\gamma) = \{i_1 < \dots < i_\ell\}$ and $x = \pi(\gamma)$.

Case 1: $r \notin M_\alpha(\gamma)$. In this case, $\delta \sim_\alpha \gamma$ implies $\delta_r = \gamma_r$. Thus it suffices to consider the case $\gamma_r = e$, as otherwise our sum is equal to zero. We can rewrite (7) as follows

$$\begin{aligned} & \sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, \delta_r = e, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} x(-\alpha_r) f'(\delta') \\ & + \sum_{\delta \in \Gamma_{s_\alpha x}, \delta \sim_\alpha \gamma, \delta_r = e, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} s_\alpha x(-\alpha_r) f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}. \end{aligned}$$

As $s_\alpha x(-\alpha_r) \equiv x(-\alpha_r) \pmod{\alpha}$, it suffices to prove that

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, \delta_r = e, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}} \quad (8)$$

and

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, \delta_r = e, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|-1}}. \quad (9)$$

We can rewrite (8) as follows

$$\sum_{\delta' \in \Gamma', \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|}}.$$

It holds by Proposition 7 applied to $f' \in \mathcal{X}'$. Noting that $J_\alpha(\delta) \subset J_\alpha(\gamma)$ is equivalent to $D_\alpha(\delta') \subset D_\alpha(\gamma')$ in (9) by Proposition 1, we can rewrite (9) as follows

$$\sum_{\delta' \in \Gamma_x, \delta' \sim_\alpha \gamma', D_\alpha(\delta') \subset D_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|-1}}.$$

By Proposition 1, we have

$$|D_\alpha(\gamma')| \geq |J_\alpha(\gamma')| - 1 = |J_\alpha(\gamma)| - 1 \quad (10)$$

and $|J_\alpha(\delta')| = |D_\alpha(\delta')|$ for $s_\alpha x > x$ and $|J_\alpha(\delta')| = |D_\alpha(\delta')| + 1$ for $s_\alpha x < x$. Therefore, the above equivalence follows from Proposition 8 applied to the element $f'|_{\Gamma'_x}$, which belongs to $\mathcal{X}'^x$ as is easy to see from the commutative diagram

$$\begin{array}{ccc} H_T^\bullet(\Sigma') & \longrightarrow & H_T^\bullet(\Sigma'_x) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma') & \longrightarrow & H_T^\bullet(\Gamma'_x) \end{array}$$

Case 2: $r \in M_\alpha(\gamma) \setminus J_\alpha(\gamma)$. In this case, $i_\ell = r$, $|D_\alpha(\gamma)| = |J_\alpha(\gamma)|$ and $s_\alpha x > x$ by Proposition 1. If δ belonged to $\Gamma_{s_\alpha x}$ in (7), we would get by Proposition 1 that $r \in J_\alpha(\delta)$ and thus the inclusion $J_\alpha(\delta) \subset J_\alpha(\gamma)$ would not hold. On the other hand, for any $\delta \in \Gamma_x$ such that $\delta \sim_\alpha \gamma$, we have $r \in M_\alpha(\delta) \setminus J_\alpha(\delta)$, whence $\beta_r(\delta) = -\alpha$. Therefore, it suffices to prove that

$$\sum_{\delta' \in \Gamma_x, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|-1}}. \quad (11)$$

If $\gamma' \in \Gamma_x$, then by Proposition 1 the summation runs over $\delta' \in \Gamma_x$ such that $\delta' \sim_\alpha \gamma'$ and $D_\alpha(\delta') \subset D_\alpha(\gamma')$. Therefore, (11) follows from (10) and Proposition 8 applied to $f'|_{\Gamma'_x}$.

We assume now that $\gamma' \in \Gamma_{xs_r} = \Gamma_{s_\alpha x}$. Note that $M_\alpha(\gamma') = \{i_1 < \dots < i_{\ell-1}\}$. This set is not empty (i.e., $\ell > 1$), as $s_\alpha \pi(\gamma') = x < s_\alpha x = \pi(\gamma')$, whence $i_{\ell-1} \in J_\alpha(\gamma')$. Consider the gallery $\tilde{\gamma}'$ that is obtained from γ' by replacing $\gamma_{i_{\ell-1}}$ with $\gamma_{i_{\ell-1}} s_{i_{\ell-1}}$. We clearly have

$$\tilde{\gamma}' \sim_\alpha \gamma', \quad \tilde{\gamma}' \in \Gamma_x, \quad J_\alpha(\tilde{\gamma}') = J_\alpha(\gamma') \setminus \{i_{\ell-1}\}, \quad D_\alpha(\tilde{\gamma}') = D_\alpha(\gamma')$$

Finally it remains to note that in (11), we have $s_\alpha \pi(\delta') = s_\alpha x > x = \pi(\delta')$, whence $i_{\ell-1} \notin J_\alpha(\delta')$. Thus $J_\alpha(\delta') \subset J_\alpha(\gamma')$ is equivalent to $J_\alpha(\delta') \subset J_\alpha(\tilde{\gamma}')$ and hence by Proposition 1 to $D_\alpha(\delta') \subset D_\alpha(\tilde{\gamma}')$. Thus we can rewrite (11) as follows

$$\sum_{\delta' \in \Gamma_x, \delta' \sim_\alpha \tilde{\gamma}', D_\alpha(\delta') \subset D_\alpha(\tilde{\gamma}')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|-1}}.$$

This equivalence again follows from (10) and Proposition 8 applied to $f'|_{\Gamma'_x}$.

Case 3: $r \in J_\alpha(\gamma)$. In this case, $i_\ell = r$ and $s_\alpha x < x$. We can rewrite (7) as follows

$$\begin{aligned} \sum_{\delta' \in \Gamma_x, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|+1} \alpha f'(\delta') \\ - \sum_{\delta' \in \Gamma_{s_\alpha x}, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} \alpha f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}. \end{aligned}$$

It suffices to prove that

$$\begin{aligned} \sum_{\delta' \in \Gamma_x, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \\ + \sum_{\delta' \in \Gamma_{s_\alpha x}, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|-1}}, \end{aligned}$$

which follows from Proposition 7, as $|J_\alpha(\gamma)| - 1 = |J_\alpha(\gamma')|$. $\square$

For notational purposes, it is convenient to define

$$\tilde{\nabla}_t f'(\gamma) = \begin{cases} \tilde{\beta}_r(\gamma) f'(\gamma') & \text{if } \gamma_r = t; \\ 0 & \text{otherwise} \end{cases}$$

Corollary 11. $\tilde{\nabla}_t f' \in \mathcal{X}$ if $f' \in \mathcal{X}'$

Proof. The result follows from $\tilde{\nabla}_e f' = \nabla_e f'$ and $\tilde{\nabla}_{s_r} f' = -\nabla_{s_r} f'$. $\square$

4.3. Folding the ends. For $r > 0$, we define the automorphism $\gamma \mapsto \dot{\gamma}$ of Γ by

$$\dot{\gamma}_i = \begin{cases} \gamma_i & \text{if } i < r; \\ s_r \gamma_r & \text{if } i = r. \end{cases}$$

It satisfies the following properties:

- $M_\alpha(\dot{\gamma}) = M_\alpha(\gamma)$;
- $\dot{\delta} \sim_\alpha \dot{\gamma} \iff \delta \sim_\alpha \gamma$;
- If $r \notin M_\alpha(\gamma)$, then $J_\alpha(\dot{\gamma}) = J_\alpha(\gamma)$. If $r \in M_\alpha(\gamma)$, then $J_\alpha(\dot{\gamma}) = J_\alpha(\gamma) \triangle \{r\}$, where $\triangle$ stands for the symmetric difference;
- $D_\alpha(\dot{\gamma}) = D_\alpha(\gamma)$;
- $\dot{\gamma} \in \Gamma_x \iff \gamma \in \Gamma_{xs_r}$,

whose proofs are left to the reader.

This automorphism of Γ induces an automorphism of $H_T^\bullet(\Gamma)$ by $\dot{f}(\gamma) = f(\dot{\gamma})$. Clearly, these automorphisms are of order 2.

Lemma 12. $\dot{\mathcal{X}} = \mathcal{X}$, $\dot{\mathcal{X}}^x = \mathcal{X}^{xs_r}$.

Proof. Actually we only have to prove that $\dot{\mathcal{X}} \subset \mathcal{X}$. Take any $f \in \mathcal{X}$. By Proposition 7, we must check the equivalence

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}} \quad (12)$$

for arbitrary $\gamma \in \Gamma$ and $\alpha \in \Phi^+$.

Case 1: $r \notin M_\alpha(\gamma)$. In this case, we can rewrite (12) as follows

$$\sum_{\delta \in \Gamma, \dot{\delta} \sim_\alpha \dot{\gamma}, J_\alpha(\dot{\delta}) \subset J_\alpha(\dot{\gamma})} (-1)^{|J_\alpha(\dot{\delta})|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|J_\alpha(\dot{\gamma})|}}.$$

It holds by Proposition 7.

Case 2: $r \in M_\alpha(\gamma) \setminus J_\alpha(\gamma)$. In this case, $\gamma \sim_\alpha \dot{\gamma}$ and $J_\alpha(\dot{\gamma}) = J_\alpha(\gamma) \sqcup \{r\}$. We can rewrite (12) as follows¹

$$\sum_{\delta \in \Gamma, \dot{\delta} \sim_\alpha \dot{\gamma}, r \in J_\alpha(\dot{\delta}), J_\alpha(\dot{\delta}) \subset J_\alpha(\dot{\gamma})} (-1)^{|J_\alpha(\dot{\delta})|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|J_\alpha(\dot{\gamma})|}}. \quad (13)$$

To prove it, let us write the following two equivalences

$$\begin{aligned} \sum_{\delta \in \Gamma, \dot{\delta} \sim_\alpha \dot{\gamma}, J_\alpha(\dot{\delta}) \subset J_\alpha(\dot{\gamma})} (-1)^{|J_\alpha(\dot{\delta})|} f(\dot{\delta}) &\equiv 0 \pmod{\alpha^{|J_\alpha(\dot{\gamma})|}}, \\ \sum_{\delta \in \Gamma, \dot{\delta} \sim_\alpha \gamma, J_\alpha(\dot{\delta}) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\dot{\delta})|} f(\dot{\delta}) &\equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}, \end{aligned}$$

which hold by Proposition 7. Subtracting the latter from the former and considering everything modulo $\alpha^{|J_\alpha(\gamma)|}$, we get (13).

Case 3: $r \in J_\alpha(\gamma)$. In this case, $\gamma \sim_\alpha \dot{\gamma}$ and we can rewrite (12) as follows²

$$- \sum_{\delta \in \Gamma, \dot{\delta} \sim_\alpha \gamma, J_\alpha(\dot{\delta}) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\dot{\delta})|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}.$$

It holds by Proposition 7. □

4.4. Fixing the ends. Let $r > 0$. Consider the natural embedding $\iota : \Sigma' \hookrightarrow \Sigma$ defined by $[p_1, \dots, p_{r-1}] \mapsto [p_1, \dots, p_{r-1}, e]$. This is a B -equivariant hence also a T -equivariant embedding. We get the following commutative diagram for restrictions:

$$\begin{array}{ccc} H_T^\bullet(\Sigma) & \longrightarrow & H_T^\bullet(\Sigma') \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma) & \longrightarrow & H_T^\bullet(\Gamma') \end{array}$$

Let f be an element of $\mathcal{X}$ (i.e., in the image of the left arrow). It follows from the commutativity of the above diagram that the composition $f' = f \circ \iota$ belongs to $\mathcal{X}'$ (i.e., to the image of the right arrow).

¹If $r \notin A$, then $B \subset A$ if and only if $r \in B \triangle \{r\}$ and $B \triangle \{r\} \subset A \triangle \{r\}$.

²If $r \in A$, then $B \subset A$ if and only if $B \triangle \{r\} \subset A$.

Lemma 13. *Let $f \in \mathcal{X}$, $r > 0$ and $t \in \{e, s_r\}$. We define $f' \in H_T^\bullet(\Gamma')$ by $f'(\gamma') = f(\gamma' \cdot t)$. Then $f' \in \mathcal{X}'$.*

Proof. The argument preceding the formulation of this lemma proves the claim for $t = e$. Now let $t = s_r$. By Lemma 12, we get $\dot{f} \in \mathcal{X}$. Then by the case $t = e$, we get $\dot{f} \circ \iota \in \mathcal{X}'$. Now the result follows from

$$\dot{f} \circ \iota(\gamma') = \dot{f}(\gamma' \cdot e) = f(\gamma' \cdot s_r) = f'(\gamma').$$

□

Lemma 14. *Let $f \in \mathcal{X}$, $r > 0$ and $t \in \{e, s_r\}$. Suppose that $f(\gamma) = 0$ for $\gamma_r \neq t$. Then $f(\gamma)$ is divisible in S by $\beta_r(\gamma)$ for any $\gamma \in \Gamma$. Moreover, the function $\gamma' \mapsto f(\gamma' \cdot t)/\beta_r(\gamma' \cdot t)$, where $\gamma' \in \Gamma'$, belongs to $\mathcal{X}'$.*

Proof. We shall prove the first claim by induction with respect to $\trianglelefteq$. So suppose that $f(\delta)$ is divisible by $\beta_r(\delta)$ for any $\delta \triangleleft \gamma$. We must prove that $f(\gamma)$ is divisible by $\beta_r(\gamma)$. Clearly, we need only to consider the case $\gamma_r = t$.

We take for α the positive of the two roots $\beta_r(\gamma)$ and $-\beta_r(\gamma)$. Thus $r \in M_\alpha(\gamma)$.

Case 1: $r \in J_\alpha(\gamma)$. In this case $|J_\alpha(\gamma)| > 0$. Thus by Proposition 7, we get

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha}.$$

As $\delta \sim_\alpha \gamma$ and $J_\alpha(\delta) \subset J_\alpha(\gamma)$ imply $\delta \trianglelefteq \gamma$, the claim follows.

Case 2: $r \notin J_\alpha(\gamma)$. In this case, $r \in J_\alpha(\dot{\gamma})$, whence $|J_\alpha(\dot{\gamma})| > 0$. Moreover, $\dot{\gamma} \sim_\alpha \gamma$. By Proposition 7, we get

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, \delta_r = t, J_\alpha(\delta) \subset J_\alpha(\dot{\gamma})} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha}. \quad (14)$$

We claim that

$$\delta \sim_\alpha \gamma, \delta_r = t, J_\alpha(\delta) \subset J_\alpha(\dot{\gamma}) \implies \delta \trianglelefteq \gamma. \quad (15)$$

We get $\delta \triangleleft \dot{\gamma}$ as $\delta_r = t \neq \dot{\gamma}_r$. Thus there exists some i_0 such that $\delta^{i_0} < \dot{\gamma}^{i_0}$ and $\delta_i = \gamma_i$ for $i < i_0$. If $i_0 < r$ then $\delta^{i_0} < \gamma^{i_0}$ and thus $\delta \triangleleft \gamma$. On the other hand, if $i_0 = r$, then $\gamma = \delta$, since $\delta_r = t = \gamma_r$.

Now it follows from (14), (15) and the inductive hypothesis that $f(\gamma)$ is divisible by α .

Let us prove the second claim. We denote by f' the function under consideration: $f'(\gamma') = f(\gamma' \cdot t)/\beta_r(\gamma' \cdot t)$. By Proposition 7, we must check the equivalence

$$\sum_{\delta' \in \Gamma', \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|}} \quad (16)$$

for any $\gamma' \in \Gamma'$ and $\alpha \in \Phi^+$. Clearly, we can assume that $J_\alpha(\gamma') \neq \emptyset$. We set $\gamma := \gamma' \cdot t$. Let us fix the notation

$$M_\alpha(\gamma) = \{i_1 < \dots < i_\ell\}, \quad y = \pi(\gamma').$$

By Proposition 7, we get

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}. \quad (17)$$

Case 1: $r \notin M_\alpha(\gamma)$. We can rewrite (17) as follows

$$\begin{aligned} yt(-\alpha_r) \sum_{\delta \in \Gamma_{yt}, \delta \sim_\alpha \gamma, \delta_r = t, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f'(\delta') \\ + s_\alpha yt(-\alpha_r) \sum_{\delta \in \Gamma_{s_\alpha yt}, \delta \sim_\alpha \gamma, \delta_r = t, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}. \end{aligned}$$

We have $s_\alpha yt(-\alpha_r) = yt(-\alpha_r) + c\alpha$ for some $c \in \mathbb{Z}$. Thus the above equivalence takes form

$$\begin{aligned} s_\alpha yt(-\alpha_r) \sum_{\delta' \in \Gamma', \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \\ - c\alpha \sum_{\delta' \in \Gamma'_y, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|}}. \end{aligned} \quad (18)$$

Our aim is to get rid of the second line. As the restriction $f|_{\Gamma_{yt}}$ belongs to $\mathcal{X}^{yt}$, Proposition 8 implies that

$$\sum_{\delta \in \Gamma_{yt}, \delta \sim_\alpha \gamma, \delta_r = t, D_\alpha(\delta) \subset D_\alpha(\gamma)} (-1)^{|D_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}},$$

which can be written as follows

$$\pm yt(-\alpha_r) \sum_{\delta' \in \Gamma_y, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}},$$

where $+$ is taken if $s_\alpha yt > yt$ and $-$ is taken otherwise (see Proposition 1). Moreover, $yt(-\alpha_r) = \gamma^r(-\alpha_r)$ is not proportional to α by the hypothesis of the current case. Hence it follows from the above equivalence that

$$c\alpha \sum_{\delta' \in \Gamma_y, \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|+1}}.$$

It remains to note that $|J_\alpha(\gamma')| = |J_\alpha(\gamma)| \geq |D_\alpha(\gamma)| + 1$, add the above equivalence to (18) and note that $s_\alpha yt(-\alpha_r)$ is also not proportional to α .

Case 2: $r \in M_\alpha(\gamma) \setminus J_\alpha(\gamma)$. In this case $i_\ell = r$, $\gamma \sim_\alpha \dot{\gamma}$, $J_\alpha(\dot{\gamma}) = J_\alpha(\gamma) \cup \{r\}$. By Proposition 7, we get

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, \delta_r = t, J_\alpha(\delta) \subset J_\alpha(\dot{\gamma})} (-1)^{|J_\alpha(\delta)|} \beta_r(\delta) f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\dot{\gamma})|}}.$$

As $r \in J_\alpha(\dot{\gamma})$, this equivalence can be rewritten as follows

$$\sum_{\delta' \in \Gamma', \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta' \cdot t)|} \beta_r(\delta' \cdot t) f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|+1}}. \quad (19)$$

Considering separately the cases $\delta' \in \Gamma'_y$ and $\delta' \in \Gamma'_{s_\alpha y}$, we get $(-1)^{|J_\alpha(\delta' \cdot t)|} \beta_r(\delta' \cdot t) f'(\delta') = (-1)^{|J_\alpha(\delta')|} yt(-\alpha_r)$. We know that $yt(-\alpha_r) = \beta_r(\gamma) = -\alpha$. Thus dividing (19) by $-\alpha$, we get (16).

Case 3: $r \in J_\alpha(\gamma)$. In this case $i_\ell = r$ and (17) can be rewritten as follows

$$\sum_{\delta' \in \Gamma', \delta' \sim_\alpha \gamma', J_\alpha(\delta') \subset J_\alpha(\gamma')} (-1)^{|J_\alpha(\delta' \cdot t)|} \beta_r(\delta' \cdot t) f'(\delta') \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma')|+1}}. \quad (20)$$

Considering separately the cases $\delta' \in \Gamma'_y$ and $\delta' \in \Gamma'_{s_\alpha y}$, we get $(-1)^{|J_\alpha(\delta' \cdot t)|} \beta_r(\delta' \cdot t) f'(\delta') = -(-1)^{|J_\alpha(\delta')|} yt(-\alpha_r)$. We know that $yt(-\alpha_r) = \beta_r(\gamma) = \alpha$. Thus dividing (20) by $-\alpha$, we get (16). $\square$

4.5. Bases for $\mathcal{X}$. For any $\rho \in \Upsilon$, we construct the subset B_ρ of $H^\bullet(\Sigma)$ inductively by

$$B_\emptyset = \{1\}, \quad B_\rho = \Delta(B_{\rho'_0}) \cup \nabla_{\rho_\emptyset}(B_{\rho'_1}).$$

By Lemmas 9 and 10, we get $B_\rho \subset \mathcal{X}$.

Theorem 15. *B_ρ is an S -basis of $\mathcal{X}$.*

Proof. We apply induction on r . This result is clearly true for $r = 0$. Therefore, we assume that $r > 0$ and that $B_{\rho'_0}$ and $B_{\rho'_1}$ are bases of $\mathcal{X}'$.

Let $f \in \mathcal{X}$. By Lemma 13, the function $f' \in H_T^\bullet(\Gamma')$ defined by $f'(\delta') = f(\delta' \cdot \rho_\emptyset s_r)$ belongs to $\mathcal{X}'$. We have

$$(f - \Delta(f'))(\delta' \cdot \rho_\emptyset s_r) = f(\delta' \cdot \rho_\emptyset s_r) - \Delta(f')(\delta' \cdot \rho_\emptyset s_r) = f'(\delta') - f'(\delta') = 0.$$

for any $\delta' \in \Gamma'$. Let us define

$$h'(\delta') = \frac{(f - \Delta(f'))(\delta' \cdot \rho_\emptyset)}{\beta_r(\delta' \cdot \rho_\emptyset)}.$$

for any $\delta' \in \Gamma'$. By Lemma 14, h' is a well defined function of $\mathcal{X}'$. The above formulas show that $f - \Delta(f') = \nabla_{\rho_\emptyset}(h')$, whence $f = \Delta(f') + \nabla_{\rho_\emptyset}(h')$. By the inductive hypothesis and the linearity of Δ and $\nabla_{\rho_\emptyset}$, the function f belongs to the S -span of B_ρ .

It remains to prove the S -linear independence of elements of B_ρ . Let $B_{\rho'_0} = \{b_1^{(0)}, \dots, b_{n_0}^{(0)}\}$ and $B_{\rho'_1} = \{b_1^{(1)}, \dots, b_{n_1}^{(1)}\}$. Suppose that we have

$$\sum_{i=1}^{n_0} \alpha_i^{(0)} \Delta(b_i^{(0)}) + \sum_{i=1}^{n_1} \alpha_i^{(1)} \nabla_{\rho_\emptyset}(b_i^{(1)}) = 0$$

for some $\alpha_i^{(0)}, \alpha_i^{(1)} \in S$. Consider the decomposition $\Gamma = \Gamma' \cdot \rho_\emptyset \sqcup \Gamma' \cdot \rho_\emptyset s_r$. Restricting the above equality to $\Gamma' \cdot \rho_\emptyset s_r$, we get $\sum_{i=1}^{n_0} \alpha_i^{(0)} b_i^{(0)} = 0$. Hence all $\alpha_i^{(0)} = 0$ and $\sum_{i=1}^{n_1} \alpha_i^{(1)} \nabla_{\rho_\emptyset}(b_i^{(1)}) = 0$. Thus $\sum_{i=1}^{n_1} \alpha_i^{(1)} \beta_r(\delta) b_i^{(1)}(\delta') = 0$ for any $\delta \in \Gamma' \cdot \rho_\emptyset s_r$. Cancelling a nonzero element $\beta_r(\delta)$, we get $\sum_{i=1}^{n_1} \alpha_i^{(1)} b_i^{(1)}(\delta') = 0$ for any $\delta' \in \Gamma'$. Hence all $\alpha_i^{(1)} = 0$. $\square$

4.6. Basis for $\mathcal{X}^x$. For any gallery $\gamma \in \Gamma$, we define

$$\mathbf{a}(\gamma) = \prod_{i \in D(\gamma)} \tilde{\beta}_i(\gamma) = \prod_{\alpha \in \Phi^+} \alpha^{|D_\alpha(\gamma)|}, \quad \mathbf{b}_\emptyset = 1, \quad \mathbf{b}_\gamma = \begin{cases} \Delta(\mathbf{b}_{\gamma'}) & \text{if } r \notin D(\gamma); \\ \tilde{\nabla}_{\gamma_r}(\mathbf{b}_{\gamma'}) & \text{if } r \in D(\gamma). \end{cases}$$

By Lemma 9 and Corollary 11, we get $\mathbf{b}_\gamma \in \mathcal{X}$.

Lemma 16. *Let $\gamma \in \Gamma_x$. Then $\mathbf{b}_\gamma(\gamma) = \mathbf{a}(\gamma)$ and $\mathbf{b}_\gamma(\delta) = 0$ for any $\delta \in \Gamma_x$ such that $\delta < \gamma$.*

Proof. The first formula follows directly from the definition of $\mathbf{b}_\gamma$ and $\tilde{\nabla}_t$. Let us prove the second claim inductively. From $\delta < \gamma$ it follows that there exists some $i_0 = 1, \dots, r-1$ such that $\delta^{i_0} < \gamma^{i_0}$ and $\delta^i = \gamma^i$ for $i > i_0$. Clearly $\delta' < \gamma'$. Assume that $\delta_r = \gamma_r$. Then $\delta', \gamma' \in \Gamma_{x\gamma_r}$. Thus by induction $\mathbf{b}_\gamma(\delta) = \mathbf{b}_{\gamma'}(\delta') = 0$ if $r \notin D(\gamma)$ and $\mathbf{b}_\gamma(\delta) = \tilde{\beta}_r(\delta) \mathbf{b}_{\gamma'}(\delta') = 0$ if $r \in D(\gamma)$. Now assume on the contrary that $\delta_r \neq \gamma_r$. However $\delta^r = \gamma^r = x$. Hence $\delta^{r-1} \neq \gamma^{r-1}$. This means that $i_0 = r-1$ and $\gamma^{r-1} s_r = \delta^{r-1} < \gamma^{r-1}$. Hence $r \in D(\gamma)$. Therefore $\mathbf{b}_\gamma(\delta) = \tilde{\nabla}_{\gamma_r}(\mathbf{b}_{\gamma'})(\delta) = 0$. $\square$

Theorem 17. *The set $\{\mathbf{b}_\gamma|_{\Gamma_x} \mid \gamma \in \Gamma_x\}$ is an S -basis of $\mathcal{X}^x$. In particular, the restriction $\mathcal{X} \rightarrow \mathcal{X}^x$ is surjective.*

Proof. This set is S -linearly independent by Lemma 16, as $<$ is the total order on Γ_x and $\mathbf{a}(\gamma) \neq 0$. Let us prove that any element $f \in \mathcal{X}^x$ is representable as an S -linear combination. We apply induction on the cardinality of the set (the upper closure of the support of f)

$$C(f) = \{\delta \in \Gamma_x \mid \text{there exists } \gamma \in \Gamma_x \text{ such that } \delta \geq \gamma \text{ and } f(\gamma) \neq 0\}.$$

If $C(f) = \emptyset$, then $f = 0$ and the result follows. Suppose now that $C(f) \neq \emptyset$ and let γ be its element minimal with respect to $<$. As

$$\delta \in \Gamma_x, \delta \sim_\alpha \gamma, D_\alpha(\delta) \subset D_\alpha(\gamma) \implies \delta \leq \gamma, \quad (21)$$

Proposition 8 implies that $f(\gamma)$ is divisible by $\prod_{\alpha \in \Phi^+} \alpha^{|D_\alpha(\gamma)|} = \mathbf{a}(\gamma)$. Consider the difference $h = f - f(\gamma)/\mathbf{a}(\gamma)\mathbf{b}_\gamma$. By Lemma 16, we get $C(h) \subset \{\delta \in \Gamma_x \mid \delta > \gamma\} \subsetneq C(f)$. By induction, h belongs to the S -span of our set. Thus so does f . $\square$

Corollary 18. *The restrictions $H_T^\bullet(\Sigma) \rightarrow H_T^\bullet(\Sigma_x)$ and $H^\bullet(\Sigma) \rightarrow H^\bullet(\Sigma_x)$ are surjective.*

Proof. The surjectivity of the first morphism follows from Theorem 17. As Σ and Σ_x are equivariantly formal, the second morphism is obtained from the first one by applying $?\otimes_S \mathbb{Z}'$ (where $S^i \mathbb{Z}' = 0$ for $i > 0$). Hence it is also surjective. $\square$

Remark 19. Let us define $\rho_r(x) \in \Upsilon$ as follows:

$$x\rho_r(x)_\emptyset > x\rho_r(x)_\emptyset s_r, \quad \rho_r(x)'_0 = \rho_{r-1}(x\rho_r(x)_\emptyset s_r), \quad \rho_r(x)'_1 = \rho_{r-1}(x\rho_r(x)_\emptyset).$$

It is elementary to check that up to sign all elements $\mathbf{b}_\gamma$ with $\gamma \in \Gamma_x$ belong to $B_{\rho_r(x)}$.

Finally, we write down the exact inductive formula for the values of the basis functions

$$\mathbf{b}_\gamma(\delta) = \begin{cases} \mathbf{b}_{\gamma'}(\delta') & \text{if } r \notin D(\gamma); \\ (\delta')^{r-1}(-\alpha_r)\mathbf{b}_{\gamma'}(\delta') & \text{if } r \in D(\gamma) \text{ and } \delta_r = \gamma_r; \\ 0 & \text{if } r \in D(\gamma) \text{ and } \delta_r \neq \gamma_r. \end{cases} \quad (22)$$

5. BASIS OF THE IMAGE $\bar{\mathcal{X}}^x$

5.1. Localization for $\bar{\Sigma}_x$. Let k be a principal ideal domain with invertible 2 if the root system contains a component of type C_n . As $\bar{\Sigma}_x$ is just a T -subspace of Σ , we can apply Theorem 5 to it as well. However, it is more difficult to apply Corollary 6. Actually, the only problem to overcome is to prove that $H_T^\bullet(\bar{\Sigma}_x, k)$ is a free S_k -module. Unfortunately, we can not solve this problem in the same way as for Σ : we do not know if $\bar{\Sigma}_x$ has an affine paving.

Consider the natural embeddings $i : \Sigma_x \hookrightarrow \Sigma$ and $j : \bar{\Sigma}_x \hookrightarrow \Sigma$. From the non-equivariant distinguished triangle

$$j_! j^* \underline{k}_\Sigma \rightarrow \underline{k}_\Sigma \rightarrow i_* i^* \underline{k}_\Sigma \xrightarrow{+1}, \quad (23)$$

we get the following exact sequence

$$H^{2m}(\Sigma, k) \rightarrow H^{2m}(\Sigma_x, k) \rightarrow H^{2m+1}(\Sigma, j_! \underline{k}_{\bar{\Sigma}_x}) \rightarrow H^{2m+1}(\Sigma, k) = 0.$$

The left morphism is surjective by Corollary 18 and the following corollary of the projection formula (cf. [I, VI.5.1]):

Proposition 20. *Let $\mathbb{Z}' \rightarrow k$ be the natural ring homomorphism. For any topological space X , we get the following exact sequence:*

$$0 \rightarrow H_c^i(X) \otimes_{\mathbb{Z}'} k \rightarrow H_c^i(X, k) \rightarrow \text{Tor}_1(H_c^{i+1}(X), k) \rightarrow 0.$$

Hence $H^{2m+1}(\Sigma, j_! \underline{k}_{\bar{\Sigma}_x}) = 0$. Since Σ is compact, we get

$$0 = H^{2m+1}(\Sigma, j_! \underline{k}_{\bar{\Sigma}_x}) = H_c^{2m+1}(\Sigma, j_! \underline{k}_{\bar{\Sigma}_x}) = H_c^{2m+1}(\bar{\Sigma}_x, k).$$

The Poincaré duality in the form [D, Theorem 3.3.3] yields the following exact sequence:
 $0 = \text{Ext}_{k\text{-mod}}^1(H_c^{2\dim \Sigma - 2m+1}(\bar{\Sigma}_x, k), k) \rightarrow H^{2m}(\bar{\Sigma}_x, k) \rightarrow \text{Hom}_{k\text{-mod}}(H_c^{2\dim \Sigma - 2m}(\bar{\Sigma}_x, k), k) \rightarrow 0.$

Hence

$$H^{2m}(\bar{\Sigma}_x, k) \simeq \text{Hom}_{k\text{-mod}}(H_c^{2\dim \Sigma - 2m}(\bar{\Sigma}_x, k), k). \quad (24)$$

From (23), we get the following exact sequence

$$0 = H^{2m-1}(\Sigma_x, k) \rightarrow H^{2m}(\Sigma, j_! \underline{k}_{\bar{\Sigma}_x}) \rightarrow H^{2m}(\Sigma, k).$$

The right cohomology is a finitely generated free k -module. We get that its submodule $H^{2m}(\Sigma, j_! \underline{k}_{\bar{\Sigma}_x}) = H_c^{2m}(\bar{\Sigma}_x, k)$ is a finitely generated free k -module. Hence and from (24), it follows that $H^{2m}(\bar{\Sigma}_x, k)$ is also a finitely generated free k -module.

Now again applying the Poincaré duality in the form [D, Theorem 3.3.3], we get

$$0 = \text{Ext}_{k\text{-mod}}^1(H_c^{2\dim \Sigma - 2m}(\bar{\Sigma}_x, k), k) \rightarrow H^{2m+1}(\bar{\Sigma}_x, k) \rightarrow \text{Hom}_{k\text{-mod}}(H_c^{2\dim \Sigma - 2m-1}(\bar{\Sigma}_x, k), k) = 0.$$

Hence we get $H^{2m+1}(\bar{\Sigma}_x, k) = 0$.

Now the degeneracy of the Leray spectral sequence at the E_2 -term implies

$$H_T^\bullet(\bar{\Sigma}_x, k) \simeq H^\bullet(\bar{\Sigma}_x) \otimes_k S_k.$$

This module is therefore a free S_k -module.

We denote by $\bar{\mathcal{X}}^x(k)$ the image of the restrictions $H_T^\bullet(\bar{\Sigma}_x, k) \rightarrow H_T^\bullet(\bar{\Gamma}_x, k)$, which is injective by the first part of Corollary 6.

5.2. Review of Härterich's constructions. We shall briefly sketch Härterich's construction, in order to be able to apply them to the cohomology of the difference $\bar{\Sigma}_x$ in Section 5.3 and prove the criterion (Proposition 21) for the image $\bar{\mathcal{X}}^x$ of the restriction $i_{\bar{\Sigma}_x, \bar{\Gamma}_x}^* : H_T^\bullet(\bar{\Sigma}_x) \rightarrow H_T^\bullet(\bar{\Gamma}_x)$.

Let α be a positive root and $\gamma \in \Gamma$. We set $T_\alpha := \ker \alpha$ and $M_\alpha(\gamma) = \{i_1 < \dots < i_\ell\}$. Härterich [H, Section 4] constructs an embedding $v_\gamma^\alpha : (G_\alpha/B_\alpha)^\ell \hookrightarrow \Sigma$ such that composition $\iota \circ v_\gamma^\alpha : (G_\alpha/B_\alpha)^\ell \rightarrow (G/B)^r$ is given by

$$(g_1, \dots, g_\ell) \mapsto (\gamma_{\min}^1, \dots, \gamma_{\min}^{i_1-1}, g_1 \gamma_{\min}^{i_1}, \dots, g_1 \gamma_{\min}^{i_2-1}, \dots, g_\ell \gamma_{\min}^{i_\ell}, \dots, g_\ell \gamma_{\min}^r),$$

where $\gamma_{\min}$ is the minimal with respect to $\triangleleft$ element in the $\sim_\alpha$ -equivalence class of γ (i.e. the unique element of this class having no load bearing α -walls). Here and in what follows we write g instead of gB_α or gB if it is clear from the context that we consider an element of G_α/B_α or G/B respectively. Note that for $\ell = 0$, the map v_γ^α takes $(G_\alpha/B_\alpha)^\ell$ isomorphically to $\{\gamma\}$.

Clearly, v_γ^α depends only on the $\sim_\alpha$ -equivalence class of γ . Corollary 4.4 from [H] states

$$\Sigma^{T_\alpha} = \bigsqcup_{\gamma \in \text{rep}(\Gamma, \sim_\alpha)} \text{im } v_\gamma^\alpha. \quad (25)$$

We note that to ensure the T -equivariance of v_γ^α , we must define an appropriate T -action on $(G_\alpha/B_\alpha)^\ell$. This can be done as follows:

$$t \cdot (g_1, \dots, g_\ell) = (tg_1 t^{-1}, \dots, tg_\ell t^{-1}).$$

Similarly the isomorphism $\iota : \Sigma_\ell^2 \xrightarrow{\sim} G_\alpha/B_\alpha$ becomes an isomorphism of T -spaces if we define the following T -action on Σ_ℓ^2 :

$$t \cdot [p_1, \dots, p_\ell] := [tp_1 t^{-1}, \dots, tp_\ell t^{-1}].$$

It is clear that $(\text{im } v_\gamma^\alpha)^T = \{\delta \in \Gamma \mid \delta \sim_\alpha \gamma\}$. We can compute the preimage of each point of the last set with respect to the map $v_\alpha^\gamma \circ \iota : \Sigma_\ell^2 \rightarrow \Sigma$. Indeed, without loss of generality, it suffices to compute $(v_\alpha^\gamma \circ \iota)^{-1}(\gamma)$. Let us define

$$g_j = \begin{cases} s_\alpha & \text{if } i_j \in J_\alpha(\gamma); \\ e & \text{otherwise.} \end{cases}$$

This definition and [H, Remark preceeding (4.1)] ensures that $v_\gamma^\alpha(g_1, \dots, g_\ell) = \gamma$. We define $\bar{g} := [g_1, g_1^{-1}g_2, \dots, g_{\ell-1}^{-1}g_\ell]$ as an element of $(G_\alpha/B_\alpha)^\ell$. Hence $v_\alpha^\gamma \circ \iota(\bar{g}) = \gamma$. The following equivalence

$$j \in J(\bar{g}) \Leftrightarrow \bar{g}^j s_\alpha < \bar{g}^j \Leftrightarrow g_j = \bar{g}^j = s_\alpha \Leftrightarrow i_j \in J_\alpha(\gamma)$$

proves that

$$i_{J(\delta)} = J_\alpha(v_\gamma^\alpha \circ \iota(\delta)) \quad (26)$$

for any $\delta \in \Gamma_\ell^2$.

Now we are going to explain how to compute the intersection $\text{im } v_\gamma^\alpha \cap \pi^{-1}(x)$. Suppose that $\ell > 0$ and $v_\gamma^\alpha(g_1, \dots, g_\ell) \in \pi^{-1}(x)$. Consider the following commutative diagram:

$$\begin{array}{ccccc} (G_\alpha/B_\alpha)^\ell & \xrightarrow{v_\gamma^\alpha} & \Sigma & \xrightarrow{\iota} & (G/B)^r \\ & & \searrow \pi & & \downarrow \text{pr}_r \\ & & & & G/B \end{array}$$

Recalling the definition of $\iota \circ v_\gamma^\alpha$ (see Härterich [H, (4.1)]), we get

$$g_\ell \gamma_{\min}^r B = xB. \quad (27)$$

If $g_\ell \in B_\alpha$, then (27) together with the Bruhat decomposition yields $x = \gamma_{\min}^r$. On the contrary, if the last equality holds, then (27) is true for any $g_\ell \in B_\alpha$.

If $g_\ell \in U_\alpha s_\alpha B_\alpha$, then $g_\ell = x_\alpha(c) s_\alpha b$ for some $c \in \mathbb{C}$ and $b \in B_\alpha$. Applying [H, (4.2)], we can rewrite (27) as follows

$$g_\ell \gamma_{\min}^r B = x_\alpha(c) s_\alpha b \gamma_{\min}^r B = x_\alpha(c) s_\alpha \gamma_{\min}^r B. \quad (28)$$

This representation is already canonical in the sense of [J, 1.13], as

$$(s_\alpha \gamma_{\min}^r)^{-1}(\alpha) = (\gamma_{\min}^r)^{-1} s_\alpha(\alpha) = (\gamma_{\min}^r)^{-1}(-\alpha) = -(\gamma_{\min}^r)^{-1}(\alpha) < 0.$$

Now comparing (28) with (27) by the Bruhat decomposition, we get $x = s_\alpha \gamma_{\min}^r$ and $c = 0$. This analysis proves the following formulas:

$$\text{im } v_\gamma^\alpha \cap \pi^{-1}(x) = \begin{cases} \emptyset & \text{if } \gamma \notin \Gamma_{\{x, s_\alpha x\}}; \\ v_\gamma^\alpha((G_\alpha/B_\alpha)^{\ell-1} \times \{e\}) & \text{if } x = \pi(\gamma_{\min}); \\ v_\gamma^\alpha((G_\alpha/B_\alpha)^{\ell-1} \times \{s_\alpha\}) & \text{if } x = s_\alpha \pi(\gamma_{\min}) \end{cases} \quad (29)$$

if $\ell > 0$ and

$$\text{im } v_\gamma^\alpha \cap \pi^{-1}(x) = \begin{cases} \emptyset & \text{if } \gamma \notin \Gamma_x; \\ \{\gamma\} & \text{if } \gamma \in \Gamma_x \end{cases} \quad (30)$$

if $\ell = 0$.

5.3. Description of $\bar{\mathcal{X}}^x$. We will prove the following analog of Propositions 7 and 8.

Proposition 21. *An element $f \in H_T^\bullet(\bar{\Gamma}_x)$ belongs to the image $\bar{\mathcal{X}}^x$ of the restriction $i_{\bar{\Sigma}_x, \bar{\Gamma}_x}^* : H_T^\bullet(\bar{\Sigma}_x) \rightarrow H_T^\bullet(\bar{\Gamma}_x)$ if and only if*

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}} \quad (31)$$

for any positive root α and gallery $\gamma \in \bar{\Gamma}_{\{x, s_\alpha x\}}$ and

$$\sum_{\delta \in \Gamma_{s_\alpha x}, \delta \sim_\alpha \gamma, D_\alpha(\delta) \subset D_\alpha(\gamma)} (-1)^{|D_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}} \quad (32)$$

for any positive root α and gallery $\gamma \in \Gamma_{s_\alpha x}$.

Proof. Subtracting $\pi^{-1}(x)$ from (25), we get

$$\bar{\Sigma}_x^{T_\alpha} = \Sigma_x^{T_\alpha} \setminus \pi^{-1}(x) = \bigsqcup_{\gamma \in \text{rep}(\Gamma, \sim_\alpha)} \text{im } v_\gamma^\alpha \setminus \pi^{-1}(x).$$

Note that $(\text{im } v_\gamma^\alpha \setminus \pi^{-1}(x))^T = \{\delta \in \bar{\Gamma}_x \mid \delta \sim_\alpha \gamma\}$.

By Corollary 4, an element $f \in H_T^\bullet(\bar{\Gamma}_x)$ belongs to the image of $H_T^\bullet(\bar{\Sigma}_x^{T_\alpha}) \rightarrow H_T^\bullet(\bar{\Gamma}_x)$ if and only if each restriction $f^\gamma := f|_{\{\delta \in \bar{\Gamma}_x \mid \delta \sim_\alpha \gamma\}}$ belongs to the image of $H_T^\bullet(\text{im } v_\gamma^\alpha \setminus \pi^{-1}(x)) \rightarrow H_T^\bullet(\{\delta \in \bar{\Gamma}_x \mid \delta \sim_\alpha \gamma\})$. Clearly, it suffices to consider only the case $\{\delta \in \bar{\Gamma}_x \mid \delta \sim_\alpha \gamma\} \neq \emptyset$. We fix such a $\gamma \in \Gamma$ and consider the set $M_\alpha(\gamma) = \{i_1 < \dots < i_\ell\}$.

Case 1: $\gamma \in \Gamma_{\{x, s_\alpha x\}}$. By (29) or (30), we get that $\text{im } v_\gamma^\alpha \setminus \pi^{-1}(x) = \text{im } v_\gamma^\alpha$ and $\{\delta \in \bar{\Gamma}_x \mid \delta \sim_\alpha \gamma\} = \{\delta \in \Gamma \mid \delta \sim_\alpha \gamma\}$. We have the commutative diagram

$$\begin{array}{ccc} \Sigma_\ell^2 & \xrightarrow[\sim]{v_\gamma^\alpha \circ \iota} & \text{im } v_\gamma^\alpha \\ \uparrow & & \uparrow \\ \Gamma_\ell^2 & \xrightarrow[\sim]{v_\gamma^\alpha \circ \iota} & (\text{im } v_\gamma^\alpha)^T = \{\delta \in \Gamma \mid \delta \sim_\alpha \gamma\} \end{array}$$

$$\begin{array}{ccccc} \Sigma_\ell^2 & \xrightarrow[\sim]{\iota} & (G_\alpha/B_\alpha)^\ell & \xrightarrow{v_\gamma^\alpha} & \text{im } v_\gamma^\alpha \\ \uparrow & & \uparrow & & \uparrow \\ \Gamma_\ell^2 & \xrightarrow[\sim]{\iota} & \{e, s_\alpha\}^\ell & \xrightarrow{v_\gamma^\alpha} & (\text{im } v_\gamma^\alpha)^T = \{\delta \in \Gamma \mid \delta \sim_\alpha \gamma\} \end{array}$$

Hence we get the following commutative diagram for cohomologies:

$$\begin{array}{ccc} H_T^\bullet(\Sigma_\ell^2) & \xleftarrow[\sim]{(v_\gamma^\alpha \circ \iota)^*} & H_T^\bullet(\text{im } v_\gamma^\alpha) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma_\ell^2) & \xleftarrow[\sim]{(v_\gamma^\alpha \circ \iota)^*} & H_T^\bullet(\{\delta \in \Gamma \mid \delta \sim_\alpha \gamma\}) \end{array}$$

Thus f^γ belongs to the image of the right vertical arrow if and only if $f \circ v_\gamma^\alpha \circ \iota$ belongs to the image of the left vertical arrow. By [H, Proposition 5.4(a)] this is equivalent to

$$\sum_{\delta \in \Gamma_\ell^2, J(\delta) \subset J(\tau)} (-1)^{|J(\delta)|} f \circ v_\gamma^\alpha \circ \iota(\delta) \equiv 0 \pmod{\alpha^{|J(\tau)|}} \quad (33)$$

for any $\tau \in \Gamma_\ell^2$. By (26), we have the following equivalences

$$|J(\delta)| = |J_\alpha(v_\gamma^\alpha \circ \iota(\delta))|, \quad J(\delta) \subset J(\tau) \Leftrightarrow J_\alpha(v_\gamma^\alpha \circ \iota(\delta)) \subset J_\alpha(v_\gamma^\alpha \circ \iota(\tau)).$$

So (33) can be rewritten as follows

$$\sum_{\delta \in \Gamma_{\ell}^2, J_{\alpha}(v_{\gamma}^{\alpha} \circ \iota(\delta)) \subset J_{\alpha}(v_{\gamma}^{\alpha} \circ \iota(\tau))} (-1)^{|J_{\alpha}(v_{\gamma}^{\alpha} \circ \iota(\delta))|} f \circ v_{\gamma}^{\alpha} \circ \iota(\delta) \equiv 0 \pmod{\alpha^{|J_{\alpha}(v_{\gamma}^{\alpha} \circ \iota(\tau))|}}.$$

Replacing $v_{\gamma}^{\alpha} \circ \iota(\delta)$ and $v_{\gamma}^{\alpha} \circ \iota(\tau)$ with δ and γ respectively, we get the final version (31).

Case 2: $x = \pi(\gamma_{\min})$. The condition $\{\delta \in \bar{\Gamma}_x \mid \delta \sim_{\alpha} \gamma\} \neq \emptyset$ implies $\ell > 0$. Let $Y = (G_{\alpha}/B_{\alpha})^{\ell-1}$ and $Z = \{e, s_{\alpha}\}^{\ell-1}$ be its set of T -fixed points. By (29), we get

$$\text{im } v_{\gamma}^{\alpha} \setminus \pi^{-1}(x) = v_{\gamma}^{\alpha}(Y \times ((G_{\alpha}/B_{\alpha}) \setminus \{e\})),$$

Consider the following maps:

- the natural inclusions $\tilde{i} : \Gamma_{\ell-1}^2 \hookrightarrow \Sigma_{\ell-1}^2$, $i_{Y,Z} : Z \hookrightarrow Y$ and $\hat{i} : \{s_{\alpha}\} \hookrightarrow (G_{\alpha}/B_{\alpha}) \setminus \{e\}$;
- $a : Y \rightarrow Y \times ((G_{\alpha}/B_{\alpha}) \setminus \{e\})$ and $b : Z \rightarrow Z \times \{s_{\alpha}\}$ that add the point s_{α} to the last position;
- the projection $p : Y \times ((G_{\alpha}/B_{\alpha}) \setminus \{e\}) \rightarrow Y$ to the first $\ell - 1$ coordinates.

By definition $p \circ a = \text{id}$. Thus $a^* \circ p^* = \text{id}$ on the level of cohomology. In particular, a^* is surjective. We have the following commutative diagram:

$$\begin{array}{ccccccc} \Sigma_{\ell-1}^2 & \xrightarrow{\sim} & Y & \xrightarrow{a} & Y \times ((G_{\alpha}/B_{\alpha}) \setminus \{e\}) & \xrightarrow{v_{\gamma}^{\alpha}} & \text{im } v_{\gamma}^{\alpha} \setminus \pi^{-1}(x) \\ \uparrow \tilde{i} & & \uparrow i_{Y,Z} & & \uparrow i_{Y,Z} \times \hat{i} & & \uparrow \\ \Gamma_{\ell-1}^2 & \xrightarrow{\sim} & Z & \xrightarrow{b} & Z \times \{s_{\alpha}\} & \xrightarrow{v_{\gamma}^{\alpha}} & (\text{im } v_{\gamma}^{\alpha} \setminus \pi^{-1}(x))^T = \{\delta \in \Gamma_{s_{\alpha}x} \mid \delta \sim_{\alpha} \gamma\} \end{array}$$

Hence we get the following commutative diagram for cohomologies:

$$\begin{array}{ccccccc} H_T^{\bullet}(\Sigma_{\ell-1}^2) & \xleftarrow{\sim} & H_T^{\bullet}(Y) & \xleftarrow{a^*} & H_T^{\bullet}(Y \times ((G_{\alpha}/B_{\alpha}) \setminus \{e\})) & \xleftarrow{(v_{\gamma}^{\alpha})^*} & H_T^{\bullet}(\text{im } v_{\gamma}^{\alpha} \setminus \pi^{-1}(x)) \\ \downarrow \tilde{i}^* & & \downarrow i_{Y,Z}^* & & \downarrow (i_{Y,Z} \times \hat{i})^* & & \downarrow \\ H_T^{\bullet}(\Gamma_{\ell-1}^2) & \xleftarrow{\sim} & H_T^{\bullet}(Z) & \xleftarrow{b^*} & H_T^{\bullet}(Z \times \{s_{\alpha}\}) & \xleftarrow{(v_{\gamma}^{\alpha})^*} & H_T^{\bullet}(\{\delta \in \Gamma_{s_{\alpha}x} \mid \delta \sim_{\alpha} \gamma\}) \end{array}$$

The surjectivity of a^* proves that b^* maps isomorphically $\text{im}(i_{Y,Z} \times \hat{i})^*$ onto $\text{im } i_{Y,Z}^*$. Therefore the same is true about the whole bottom line of the above diagram: the image of the rightmost vertical arrow is mapped isomorphically onto $\text{im } \tilde{i}^*$. Thus f^{γ} belongs to the image of the rightmost vertical arrow if and only if $f \circ v_{\gamma}^{\alpha} \circ b \circ \iota \in \text{im } \tilde{i}^*$. By [H, Proposition 5.4(a)] this is equivalent to

$$\sum_{\delta \in \Gamma_{\ell-1}^2, J(\delta) \subset J(\tau)} (-1)^{|J(\delta)|} f \circ v_{\gamma}^{\alpha} \circ b \circ \iota(\delta) \equiv 0 \pmod{\alpha^{|J(\tau)|}}. \quad (34)$$

for any $\tau \in \Gamma_{\ell-1}^2$. Consider the commutative diagram

$$\begin{array}{ccc} \Gamma_{\ell-1}^2 & \xrightarrow{\iota} & Z \\ q \downarrow & & \downarrow b \\ \Gamma_{\ell}^2 & \xrightarrow{\iota} & Z \times \{s_{\alpha}\} \end{array}$$

where $q(\delta_1, \dots, \delta_{\ell-1}) = (\delta_1, \dots, \delta_{\ell-1}, \delta_{\ell-1} \cdots \delta_2 \delta_1 s_{\alpha})$. Note that $v_{\gamma}^{\alpha} \circ b \circ \iota(\delta)$ runs over the set $\{\delta \in \Gamma_{s_{\alpha}x} \mid \delta \sim_{\alpha} \gamma\}$ as δ runs over $\Gamma_{\ell-1}^2$. By (26), we get

$$J_{\alpha}(v_{\gamma}^{\alpha} \circ b \circ \iota(\delta)) = J_{\alpha}(v_{\gamma}^{\alpha} \circ \iota \circ q(\delta)) = i_{J(q(\delta))} = i_{J(\delta)} \sqcup \{i_{\ell}\}.$$

Hence

$$|D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\delta))| = |J_\alpha(v_\gamma^\alpha \circ b \circ \iota(\delta)) \setminus \{i_\ell\}| = |i_{J(\delta)}| = |J(\delta)|.$$

By (26) and Proposition 1, the inclusion relation is also preserved:

$$J(\delta) \subset J(\tau) \Leftrightarrow J_\alpha(v_\gamma^\alpha \circ b \circ \iota(\delta)) \subset J_\alpha(v_\gamma^\alpha \circ b \circ \iota(\tau)) \Leftrightarrow D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\delta)) \subset D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\tau)).$$

Now we can replace (34) with

$$\sum_{\delta \in \Gamma_{\ell-1}^2, D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\delta)) \subset D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\tau))} (-1)^{|D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\delta))|} f \circ v_\gamma^\alpha \circ b \circ \iota(\delta) \equiv 0 \pmod{\alpha^{|D_\alpha(v_\gamma^\alpha \circ b \circ \iota(\tau))|}}.$$

Replacing $v_\gamma^\alpha \circ b \circ \iota(\delta)$ and $v_\gamma^\alpha \circ b \circ \iota(\tau)$ with δ and γ respectively, we get (32).

Case 3: $x = s_\alpha \pi(\gamma_{\min})$. For $\ell > 0$, this case can be obtained from Case 2 by interchanging e and s_α . We get the following version of (34):

$$\sum_{\delta \in \Gamma_{\ell-1}^2, J(\delta) \subset J(\tau)} (-1)^{|J(\delta)|} f \circ v_\gamma^\alpha \circ b' \circ \iota(\delta) \equiv 0 \pmod{\alpha^{|J(\tau)|}}. \quad (35)$$

where $b' : Z \rightarrow Z \times \{e\}$ adds the point e to the last position. We have a similar commutative diagram

$$\begin{array}{ccc} \Gamma_{\ell-1}^2 & \xrightarrow{\iota} & Z \\ q' \downarrow & & \downarrow b' \\ \Gamma_\ell^2 & \xrightarrow{\iota} & Z \times \{e\} \end{array}$$

where $q'(\delta_1, \dots, \delta_{\ell-1}) = (\delta_1, \dots, \delta_{\ell-1}, \delta_{\ell-1} \cdots \delta_2 \delta_1)$. Here again $v_\gamma^\alpha \circ b \circ \iota(\delta)$ runs over the set $\{\delta \in \Gamma_{s_\alpha x} \mid \delta \sim_\alpha \gamma\}$ as δ runs over $\Gamma_{\ell-1}^2$. By (26), we get

$$J_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\delta)) = J_\alpha(v_\gamma^\alpha \circ \iota \circ q'(\delta)) = i_{J(q'(\delta))} = i_{J(\delta)}.$$

Hence

$$|D_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\delta))| = |J_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\delta)) \setminus \{i_\ell\}| = |i_{J(\delta)}| = |J(\delta)|$$

and the same is true for the inclusion

$$J(\delta) \subset J(\tau) \Leftrightarrow J_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\delta)) \subset J_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\tau)) \Leftrightarrow D_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\delta)) \subset D_\alpha(v_\gamma^\alpha \circ b' \circ \iota(\tau)).$$

Therefore, we again get (32).

Finally, assume that $\ell = 0$. In this case, $\gamma = \gamma_{\min}$ and $\text{im } v_\gamma^\alpha \setminus \pi^{-1}(x) = \{\gamma\}$. Hence the restriction $H_T^\bullet(\text{im } v_\gamma^\alpha \setminus \pi^{-1}(x)) \rightarrow H_T^\bullet(\{\delta \in \bar{\Gamma}_x \mid \delta \sim_\alpha \gamma\})$ is the identity map. Thus f^γ is always in its image. On the other hand, condition (32) is also satisfied, as $D_\alpha(\gamma) = \emptyset$.

All the cases being considered, it suffices to apply Corollary 6 in order to conclude the proof. $\square$

5.4. Versions of Lemmas 12, 13, and 14 for $\bar{\Sigma}_x$. First, we get the following results similar to the first two lemmas.

Lemma 22 (cf. Lemma 12). $\dot{\mathcal{X}}^x = \bar{\mathcal{X}}^{xs_r}$.

Proof. Take any $f \in \bar{\mathcal{X}}^x$. By Proposition 21, in order to prove that $\dot{f} \in \bar{\mathcal{X}}^{xs_r}$, we must check the following equivalences:

$$\sum_{\delta \in \Gamma, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}$$

for $\gamma \in \bar{\Gamma}_{\{xs_r, s_\alpha xs_r\}}$ and

$$\sum_{\delta \in \Gamma_{s_\alpha xs_r}, \delta \sim_\alpha \gamma, D_\alpha(\delta) \subset D_\alpha(\gamma)} (-1)^{|D_\alpha(\delta)|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}}$$

for $\gamma \in \Gamma_{s_\alpha x s_r}$. The first equivalence can be proved exactly as in Lemma 12. Note that $\gamma, \dot{\gamma} \in \bar{\Gamma}_{\{x, s_\alpha x\}}$ if $r \in M_\alpha(\gamma)$ (for Case 2). In view of the properties listed in Section 4.3, the second equivalence can be rewritten in the form

$$\sum_{\dot{\delta} \in \Gamma_{s_\alpha x}, \dot{\delta} \sim_\alpha \dot{\gamma}, D_\alpha(\dot{\delta}) \subset D_\alpha(\dot{\gamma})} (-1)^{|D_\alpha(\dot{\delta})|} f(\dot{\delta}) \equiv 0 \pmod{\alpha^{|D_\alpha(\dot{\gamma})|}}.$$

It holds by Proposition 21. $\square$

Lemma 23 (cf. Lemma 13). *Let $f \in \bar{\mathcal{X}}^x$, $r > 0$ and $t \in \{e, s_r\}$. We define $f' \in H_T^\bullet(\bar{\Gamma}'_{xt})$ by $f'(\gamma') = f(\gamma' \cdot t)$. Then $f' \in (\bar{\mathcal{X}}')^{xt}$.*

Proof. The same embedding ι as in Section 4.4 induces the commutative diagram

$$\begin{array}{ccc} H_T^\bullet(\bar{\Sigma}_x) & \longrightarrow & H_T^\bullet(\bar{\Sigma}'_x) \\ \downarrow & & \downarrow \\ H_T^\bullet(\bar{\Gamma}_x) & \longrightarrow & H_T^\bullet(\bar{\Gamma}'_x) \end{array}$$

Therefore the lemma holds for $t = e$. In order to prove it for $t = s_r$, consider $\dot{f}$ and apply Lemma 22. $\square$

The version of Lemma 14 requires however a more accurate choice of t .

Lemma 24 (cf. Lemma 14). *Let $x \in W$, $f \in \bar{\mathcal{X}}^x$ and $r > 0$. We can choose a unique $t \in \{e, s_r\}$ such that $xt < xts_r$. Suppose that $f(\gamma) = 0$ for $\gamma_r \neq t$. Then $f(\gamma)$ is divisible by $\beta_r(\gamma)$ for any $\gamma \in \bar{\Gamma}_x$. Moreover, the function $\gamma' \mapsto f(\gamma' \cdot t)/\beta_r(\gamma' \cdot t)$, where $\gamma' \in \bar{\Gamma}'_{xt}$, belongs to $(\bar{\mathcal{X}}')^{xt}$.*

Proof. First, we show how to choose t . Let us choose $t \in \{e, s_r\}$ arbitrarily. The elements xt and xts_r are comparable with respect to the Bruhat order, as they differ by a reflection. The element t is already chosen if $xt < xts_r$. Suppose that $xt > xts_r$. Then we set $t' = ts_r$ and get $xt' = xts_r < xt = xt's_r$. The uniqueness is clear from $xt < xts_r \Leftrightarrow xt' > xt's_r$ with t' as before.

As in the proof of Lemma 14, we shall prove the divisibility claim by induction with respect to $\trianglelefteq$. So suppose that $f(\delta)$ is divisible by $\beta_r(\delta)$ for any $\delta \in \bar{\Gamma}_x$ such that $\delta \triangleleft \gamma$ for some $\gamma \in \bar{\Gamma}_x$. We must prove that $f(\gamma)$ is divisible by $\beta_r(\gamma)$. Clearly, we need only to consider the case $\gamma_r = t$.

We take for α the positive of the two roots $\beta_r(\gamma)$ and $-\beta_r(\gamma)$. Thus $r \in M_\alpha(\gamma)$. Note the following chain of equivalences:

$$\gamma \in \Gamma_{s_\alpha x} \Leftrightarrow \gamma^r = s_\alpha x = \gamma^r s_r (\gamma^r)^{-1} x \Leftrightarrow \gamma^r s_r = x \Leftrightarrow \gamma^r = x s_r \Leftrightarrow \gamma \in \Gamma_{x s_r}. \quad (36)$$

Similarly, we get $\dot{\gamma} \in \Gamma_{s_\alpha x} \Leftrightarrow \dot{\gamma} \in \Gamma_{x s_r}$.

The case $\gamma \in \bar{\Gamma}_{\{x, s_\alpha x\}}$ is identical to Cases 1 and 2 of Lemma 14, where one applies Proposition 21 instead of Proposition 7. Note that in this case $\dot{\gamma} \in \bar{\Gamma}_{\{x, s_\alpha x\}}$ by (36), Case 1 corresponds to $r \in J_\alpha(\gamma)$ and Case 2 to $r \notin J_\alpha(\gamma)$.

Consider the case $\gamma \in \Gamma_{s_\alpha x}$. By Proposition 21, we get

$$\sum_{\delta \in \Gamma_{s_\alpha x}, \delta \sim_\alpha \gamma, D_\alpha(\delta) \subset D_\alpha(\gamma)} (-1)^{|D_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}}.$$

We have $\pi(\delta) = \pi(\gamma)$ in the summation. It follows from this fact and Proposition 1 that $J_\alpha(\delta) \subset J_\alpha(\gamma)$. Hence $\delta \trianglelefteq \gamma$.

It remains to check that $D_\alpha(\gamma) \neq \emptyset$. By (36), we get $\gamma \in \Gamma_{x s_r}$. Thus $\gamma^{r-1} = \gamma^r t = x s_r t$, whence our condition $xt < xts_r$ implies $\gamma^{r-1} s_r < \gamma^{r-1}$ and $r \in D_\alpha(\gamma)$.

Let us prove the last claim. We denote by f' the function under consideration: $f'(\gamma') = f(\gamma' \cdot t) / \beta_r(\gamma' \cdot t)$. By Proposition 21, we must check the equivalence

$$\sum_{\delta' \in \Gamma', \delta' \sim_{\alpha} \gamma', J_{\alpha}(\delta') \subset J_{\alpha}(\gamma')} (-1)^{|J_{\alpha}(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|J_{\alpha}(\gamma')|}}$$

for any $\gamma' \in \bar{\Gamma}'_{\{xt, s_{\alpha}xt\}}$ and the equivalence

$$\sum_{\delta' \in \Gamma'_{s_{\alpha}xt}, \delta' \sim_{\alpha} \gamma', D_{\alpha}(\delta') \subset D_{\alpha}(\gamma')} (-1)^{|D_{\alpha}(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|D_{\alpha}(\gamma')|}} \quad (37)$$

for any $\gamma' \in \bar{\Gamma}'_{s_{\alpha}xt}$. The first one can be proved exactly as in Lemma 14, as $\gamma \in \bar{\Gamma}_{\{x, s_{\alpha}x\}}$ and $\dot{\gamma} \in \bar{\Gamma}_{\{x, s_{\alpha}x\}}$ if $r \in M_{\alpha}(\gamma)$ (Case 2), where $\gamma = \gamma' \cdot t$.

It remains to prove (37). In this case, $\gamma = \gamma' \cdot t \in \Gamma_{s_{\alpha}x}$. We consider the following cases.

Case a: $r \notin M_{\alpha}(\gamma)$. In this case $\beta_t(\gamma) \neq \pm\alpha$. Hence $s_{\alpha}x(-\alpha_r) \neq \pm\alpha$. By Proposition 21, we get

$$\sum_{\delta \in \Gamma_{s_{\alpha}x}, \delta \sim_{\alpha} \gamma, \delta_r = t, D_{\alpha}(\delta) \subset D_{\alpha}(\gamma)} (-1)^{|D_{\alpha}(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|D_{\alpha}(\gamma)|}}. \quad (38)$$

As $\delta_r = \gamma_r = t$ in this summation, we can rewrite the above equivalence as follows

$$s_{\alpha}x(-\alpha_r) \sum_{\delta' \in \Gamma_{s_{\alpha}xt}, \delta' \sim_{\alpha} \gamma', D_{\alpha}(\delta') \subset D_{\alpha}(\gamma')} (-1)^{|D_{\alpha}(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|D_{\alpha}(\gamma')|}}.$$

Cancelling out $s_{\alpha}x(-\alpha_r)$, we get (37).

Case b: $r \in M_{\alpha}(\gamma)$. In this case, $xs_r = s_{\alpha}x$. For any $\delta \in \Gamma_{s_{\alpha}x}$ such that $\delta_r = t$ and $\delta \sim_{\alpha} \gamma$, we have $\delta^{r-1}s_r = s_{\alpha}xts_r = xt < xts_r = s_{\alpha}xt = \delta^{r-1}$. Hence $r \in D_{\alpha}(\delta)$. Therefore we can rewrite (38) as follows

$$\pm\alpha \sum_{\delta' \in \Gamma_{s_{\alpha}xt}, \delta' \sim_{\alpha} \gamma', D_{\alpha}(\delta') \subset D_{\alpha}(\gamma')} (-1)^{|D_{\alpha}(\delta')|} f'(\delta') \equiv 0 \pmod{\alpha^{|D_{\alpha}(\gamma')|+1}}$$

Cancelling out $\pm\alpha$, we get (37). □

5.5. Basis for $\bar{\mathcal{X}}^x$. For any gallery $\gamma \in \Gamma$ and $x \in W$, we define

$$\mathbf{c}_{\emptyset}^x = 1, \quad \mathbf{c}_{\gamma}^x = \begin{cases} \Delta(\mathbf{c}_{\gamma'}^{x\gamma_r}) & \text{if } x\gamma_r > x\gamma_r s_r; \\ \nabla_{\gamma_r}(\mathbf{c}_{\gamma'}^{x\gamma_r}) & \text{if } x\gamma_r < x\gamma_r s_r. \end{cases}$$

By Lemmas 9 and 10, we get $\mathbf{c}_{\gamma}^x \in \mathcal{X}$.

Theorem 25. *The set $\{\mathbf{c}_{\gamma}^x|_{\bar{\Gamma}_x} \mid \gamma \in \bar{\Gamma}_x\}$ is an S -basis of $\bar{\mathcal{X}}^x$. In particular, the restrictions $\mathcal{X} \rightarrow \bar{\mathcal{X}}^x$ and $H_T^{\bullet}(\Sigma) \rightarrow H_T^{\bullet}(\bar{\Sigma}_x)$ are surjective.*

Proof. We apply induction on r , the result being obvious for $r = 0$. Now let $r > 0$ and f be an element of $\bar{\mathcal{X}}^x$. Choose $q \in \{e, s_r\}$ so that $xq > xqs_r$ and define $f'(\gamma') = f(\gamma' \cdot q)$ for $\gamma' \in \bar{\Gamma}'_{xq}$. By Lemma 23, we get $f' \in (\bar{\mathcal{X}}')^{xq}$. By the inductive hypothesis, $f' = \sum_{\gamma' \in \bar{\Gamma}'_{xq}} a_{\gamma'} \mathbf{c}_{\gamma'}^{xq}|_{\bar{\Gamma}'_{xq}}$ for some $a_{\gamma'} \in S$. Consider the difference

$$h = f - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = q} a_{\gamma'} \mathbf{c}_{\gamma}^x|_{\bar{\Gamma}_x}. \quad (39)$$

By the above definitions, we get $h(\delta) = 0$ for any $\delta \in \bar{\Gamma}_x$ such that $\delta_r = q$:

$$\begin{aligned} h(\delta) &= f(\delta) - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = q} a_{\gamma'} \mathbf{c}_{\gamma}^x(\delta) = f'(\delta') - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = q} a_{\gamma'} \Delta(\mathbf{c}_{\gamma'}^{xq})(\delta) = \\ &= \sum_{\gamma' \in \bar{\Gamma}'_{xq}} a_{\gamma'} \mathbf{c}_{\gamma'}^{xq}(\delta') - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = q} a_{\gamma'} \mathbf{c}_{\gamma'}^{xq}(\delta') = 0. \end{aligned}$$

Let t be the element of $\{e, s_r\}$ distinct from q . We clearly have $xt < xts_r$. Thus by Lemma 24, we get that the function h' defined by $h'(\gamma') = h(\gamma' \cdot t) / \beta_r(\gamma' \cdot t)$ for $\gamma' \in \bar{\Gamma}'_{xt}$ is a well-defined element of $(\mathcal{X}')^{xt}$. By induction, $h' = \sum_{\gamma' \in \bar{\Gamma}'_{xt}} b_{\gamma'} \mathbf{c}_{\gamma'}^{xt} |_{\bar{\Gamma}'_{xt}}$ for some $b_{\gamma'} \in S$. We get $h = \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} b_{\gamma'} \mathbf{c}_{\gamma}^x |_{\bar{\Gamma}_x}$. Indeed both sides evaluate to 0 at $\delta \in \bar{\Gamma}_x$ such that $\delta_r = q$ and for $\delta \in \bar{\Gamma}_x$ such that $\delta_r = t$, we get

$$\begin{aligned} h(\delta) - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} b_{\gamma'} \mathbf{c}_{\gamma}^x(\delta) &= h(\delta' \cdot t) - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} b_{\gamma'} \nabla_t(\mathbf{c}_{\gamma'}^{xt})(\delta' \cdot t) \\ &= \beta_r(\delta) h'(\delta') - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} b_{\gamma'} \beta_r(\delta) \mathbf{c}_{\gamma'}^{xt}(\delta') = \\ &= \beta_r(\delta) \sum_{\gamma' \in \bar{\Gamma}'_{xt}} b_{\gamma'} \mathbf{c}_{\gamma'}^{xt}(\delta') - \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} b_{\gamma'} \beta_r(\delta) \mathbf{c}_{\gamma'}^{xt}(\delta') = 0. \end{aligned}$$

Hence and from (39), we get that f is a S -linear combination of elements of our set.

Finally, let us prove the linear independence. Suppose that we have $\sum_{\gamma \in \bar{\Gamma}_x} a_{\gamma} \mathbf{c}_{\gamma}^x = 0$ for some $a_{\gamma} \in S$. We can write this sum as follows

$$\sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = q} a_{\gamma} \Delta(\mathbf{c}_{\gamma'}^{xq}) + \sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} a_{\gamma} \nabla_t(\mathbf{c}_{\gamma'}^{xt}) = 0. \quad (40)$$

Evaluation at $\delta \in \bar{\Gamma}_x$ with $\delta_r = q$ yields $\sum_{\gamma' \in \bar{\Gamma}'_{xq}} a_{\gamma} \mathbf{c}_{\gamma'}^{xq}(\delta') = 0$. Hence by the inductive hypothesis, $a_{\gamma} = 0$ for any $\gamma \in \bar{\Gamma}_x$ such that $\gamma_r = q$. Therefore (40) takes the form $\sum_{\gamma \in \bar{\Gamma}_x, \gamma_r = t} a_{\gamma} \nabla_t(\mathbf{c}_{\gamma'}^{xt}) = 0$. Evaluation at $\delta \in \bar{\Gamma}_x$ with $\delta_r = t$ yields $\sum_{\gamma' \in \bar{\Gamma}'_{xt}} a_{\gamma} \beta_r(\delta) \mathbf{c}_{\gamma'}^{xt}(\delta') = 0$, whence $\sum_{\gamma' \in \bar{\Gamma}'_{xt}} a_{\gamma} \mathbf{c}_{\gamma'}^{xt}(\delta') = 0$. By the inductive hypothesis $a_{\gamma} = 0$ for any $\gamma \in \bar{\Gamma}_x$ such that $\gamma_r = t$. $\square$

Remark 26. We have $\{\mathbf{c}_{\gamma}^x \mid \gamma \in \Gamma\} = B_{\xi_r(x)}$, where $\xi_r(x) \in \Upsilon$ is defined inductively as follows: $\xi_0(x) = e$ and

$$x\xi_r(x)_{\emptyset} < x\xi_r(x)_{\emptyset} s_r, \quad \xi_r(x)'_0 = \xi_{r-1}(x\xi_r(x)_{\emptyset} s_r), \quad \xi_r(x)'_1 = \xi_{r-1}(x\xi_r(x)_{\emptyset}).$$

6. THE COSTALK-TO-STALK EMBEDDING AND THE DECOMPOSITION OF THE DIRECT IMAGE

6.1. Change of coefficients. Let k be a principal ideal domain with invertible 2 if the root system contains a component of type C_n . Consider the canonical ring homomorphism $\mathbb{Z}' \rightarrow k$. It extends to the ring homomorphism $S \rightarrow S_k$. We get the following commutative diagram:

$$\begin{array}{ccc} H^i(\Sigma) \otimes_{\mathbb{Z}'} k & \longrightarrow & H^i(\Sigma, k) \\ \downarrow & & \downarrow \\ H^i(\Gamma) \otimes_{\mathbb{Z}'} k & \longrightarrow & H^i(\Gamma, k) \end{array} \quad (41)$$

As $H^i(\Sigma)$ and $H^i(\Sigma, k)$ vanish in odd degrees, Σ is compact and Γ is finite, Proposition 20 implies that the horizontal arrows are isomorphisms. Hence we get the following chain of isomorphisms:

$$H_T^\bullet(\Sigma) \otimes_S S_k \simeq (H^\bullet(\Sigma) \otimes_{\mathbb{Z}'} S) \otimes_S S_k \simeq (H^\bullet(\Sigma) \otimes_{\mathbb{Z}'} k) \otimes_k S_k \simeq H^\bullet(\Sigma, k) \otimes_k S_k \simeq H_T^\bullet(\Sigma, k).$$

The similar chain yields an isomorphism $H_T^\bullet(\Gamma) \otimes_S S_k \xrightarrow{\sim} H_T^\bullet(\Gamma, k)$. Diagram (41) proves that these isomorphisms are compatible with the restriction from Σ to Γ . This means that we get the following commutative diagram:

$$\begin{array}{ccccc} & & H_T^\bullet(\Sigma) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Sigma, k) \\ & \nearrow \sim & \downarrow & & \downarrow \\ \mathcal{X} \otimes_S S_k & \hookrightarrow & H_T^\bullet(\Gamma) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Gamma, k) \end{array}$$

If we go along the upper path, then we get an isomorphism of S_k -modules $\mathcal{X} \otimes_S S_k \xrightarrow{\sim} \mathcal{X}(k)$. However this map is the same as the map of the lower path. Hence we get the following result.

Lemma 27. *There exist an isomorphism of S_k -modules (dashed arrow) such that the following diagram is commutative:*

$$\begin{array}{ccc} \mathcal{X} \otimes_S S_k & \xrightarrow[\sim]{\exists} & \mathcal{X}(k) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Gamma, k) \end{array}$$

Arguing similarly with Σ_x and Γ_x , we get the following results.

Lemma 28. *There exist an isomorphism of S_k -modules (dashed arrow) such that the following diagram is commutative:*

$$\begin{array}{ccc} \mathcal{X}^x \otimes_S S_k & \xrightarrow[\sim]{\exists} & \mathcal{X}^x(k) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma_x) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Gamma_x, k) \end{array}$$

The case of $\mathcal{X}^x(k)$ is more difficult, as $\bar{\Sigma}_x$ is in general not compact. However, we can use the Poincaré duality $H^i(\bar{\Sigma}_x, k) \simeq \text{Hom}_{k\text{-mod}}(H_c^{2\dim \Sigma - i}(\bar{\Sigma}_x, k), k)$ establishes in Section 5.1. We get the following sequence of canonical maps:

$$\begin{aligned} H^i(\bar{\Sigma}_x) \otimes_{\mathbb{Z}'} k &\simeq \text{Hom}_{\mathbb{Z}'\text{-mod}}(H_c^{2\dim \Sigma - i}(\bar{\Sigma}_x), \mathbb{Z}') \otimes_{\mathbb{Z}'} k \xrightarrow{\varphi} \text{Hom}_{k\text{-mod}}(H_c^{2\dim \Sigma - i}(\bar{\Sigma}_x) \otimes_{\mathbb{Z}'} k, k) \\ &\simeq \text{Hom}_{k\text{-mod}}(H_c^{2\dim \Sigma - i}(\bar{\Sigma}_x, k), k) \simeq H^i(\bar{\Sigma}_x, k). \end{aligned}$$

From Section 5.1, we know that $H_c^{2\dim \Sigma - i}(\bar{\Sigma}_x)$ is a finitely generated free $\mathbb{Z}'$ -module. Hence we conclude that the morphism φ in the sequence above is an isomorphism. If we replace $\bar{\Sigma}_x$ by $\bar{\Gamma}_x$ in this argument, then we get an isomorphism $H^i(\bar{\Gamma}_x) \otimes_{\mathbb{Z}'} k \simeq H^i(\bar{\Gamma}_x, k)$. It is rather easy to see that these isomorphisms are compatible with the restriction from $\bar{\Sigma}_x$ to $\bar{\Gamma}_x$. An argument similar to the one preceding Lemma 27, proves the following result.

Lemma 29. *There exist an isomorphism of S_k -modules (dashed arrow) such that the following diagram is commutative:*

$$\begin{array}{ccc} \bar{\mathcal{X}}^x \otimes_S S_k & \overset{\exists}{\underset{\sim}{\dashrightarrow}} & \bar{\mathcal{X}}^x(k) \\ \downarrow & & \downarrow \\ H_T^\bullet(\bar{\Gamma}_x) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\bar{\Gamma}_x, k) \end{array}$$

These three lemmas allow us to construct bases of $\mathcal{X}(k)$, $\mathcal{X}^x(k)$, $\bar{\mathcal{X}}(k)^x$ from the bases of $\mathcal{X}$, $\mathcal{X}^x$, $\bar{\mathcal{X}}^x$ given by Theorems 15, 17, 25 respectively. Moreover, we can construct operators Δ and ∇_t on $H_T^\bullet(\Gamma', k)$ similarly to Section 4.2 and obtain analogs of Lemmas 9 and 10.

6.2. Description of the costalk-to-stalk embedding. From the T -equivariant distinguished triangle

$$i_* i^! \underline{k}_\Sigma \rightarrow \underline{k}_\Sigma \rightarrow j_* j^* \underline{k}_\Sigma \xrightarrow{+1},$$

where i and j are as in Section 5.1, we get the following exact sequence

$$H_T^n(\Sigma_x, i^! \underline{k}_\Sigma) \rightarrow H_T^n(\Sigma, k) \rightarrow H_T^n(\bar{\Sigma}_x, k) \rightarrow H_T^{n+1}(\Sigma_x, i^! \underline{k}_\Sigma) \rightarrow \quad (42)$$

We are actually interested in the left map. It would be very convenient if we could prove that its source $H_T^n(\Sigma_x, i^! \underline{k}_\Sigma)$ vanishes in odd degrees. This is fortunately true, as the sequence

$$H_T^{2m}(\Sigma, k) \rightarrow H_T^{2m}(\bar{\Sigma}_x, k) \rightarrow H_T^{2m+1}(\Sigma_x, i^! \underline{k}_\Sigma) \rightarrow H_T^{2m+1}(\Sigma, k) = 0$$

is exact by (42) and the left map is surjective by Theorem 25 and Lemmas 27 and 29.

Consider the following commutative diagram with the exact first row:

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_T^{2m}(\Sigma_x, i^! \underline{k}_\Sigma) & \longrightarrow & H_T^{2m}(\Sigma, k) & \longrightarrow & H_T^{2m}(\bar{\Sigma}_x, k) \longrightarrow 0 \\ & & \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ 0 & \longrightarrow & \ker \varphi^{2m} & \longrightarrow & \mathcal{X}(k)^{2m} & \xrightarrow{\varphi^{2m}} & \bar{\mathcal{X}}^x(k)^{2m} \longrightarrow 0 \end{array}$$

Here φ^{2m} is the restriction map $f \mapsto f|_{\bar{\Gamma}_x}$ and the solid vertical arrows are induced by embeddings $\Gamma \hookrightarrow \Sigma$ and $\bar{\Gamma}_x \hookrightarrow \bar{\Sigma}_x$ respectively. We thus have proved the following result.

Lemma 30. *There exists an isomorphism of S_k -modules (dashed arrow) such that the following diagram is commutative:*

$$\begin{array}{ccc} H_T^\bullet(\Sigma_x, i^! \underline{k}_\Sigma) & \longrightarrow & H_T^\bullet(\Sigma, k) \\ \downarrow \wr & & \downarrow \wr \\ \mathcal{X}_x(k) & \hookrightarrow & \mathcal{X}(k) \end{array}$$

where $\mathcal{X}_x(k) = \{f \in \mathcal{X}(k) \mid f|_{\bar{\Gamma}_x} = 0\}$ and the bottom arrow is the natural embedding.

Corollary 31. *The functor $H^\bullet(\Sigma_x, -)$ applied to the natural morphism $i^! \underline{k}_\Sigma \rightarrow i^* \underline{k}_\Sigma$, where $i : \Sigma_x \hookrightarrow \Sigma$ is the embedding, yields a map isomorphic to the embedding $\mathcal{X}_x(k) \hookrightarrow \mathcal{X}^x(k)$.*

It remains to discuss the behavior of $\mathcal{X}_x(k)$ with respect to the change of the ring of coefficients k . By the remark at the end of Section 5.5, we have $\{\mathbf{c}_\gamma^x \mid \gamma \in \Gamma\} = B_{\xi_r(x)}$. Thus we can write $B_{\xi_r(x)} = \{b_1, \dots, b_m, b_{m+1}, \dots, b_n\}$ so that $\{b_1|_{\bar{\Gamma}_x}, \dots, b_m|_{\bar{\Gamma}_x}\}$ is a basis of

$\bar{\mathcal{X}}^x$. We have the following decompositions $b_j|_{\bar{\Gamma}_x} = \sum_{i=1}^m c_{i,j} b_i|_{\bar{\Gamma}_x}$ for some (homogeneous) $c_{i,j} \in S$. Let $u = \sum_{i=1}^n x_i b_i$, where $x_i \in S$, be an arbitrary element of $\mathcal{X}$. We get

$$u|_{\bar{\Gamma}_x} = \sum_{i=1}^n x_i b_i|_{\bar{\Gamma}_x} = \sum_{i=1}^m x_i b_i|_{\bar{\Gamma}_x} + \sum_{j=m+1}^n x_j \sum_{i=1}^m c_{i,j} b_i|_{\bar{\Gamma}_x} = \sum_{i=1}^m \left(x_i + \sum_{j=m+1}^n c_{i,j} x_j \right) b_i|_{\bar{\Gamma}_x}.$$

Hence $\mathcal{X}_x = \mathcal{X}_x(\mathbb{Z}')$ is a free S -module with basis $\left\{ -\sum_{i=1}^m c_{i,j} b_i + b_j \right\}_{j=m+1}^n$. Arguing similarly, we get that $\mathcal{X}_x(k)$ is a free S_k -module with basis $\left\{ -\sum_{i=1}^m (c_{i,j} \otimes 1_k)(b_i \otimes 1_k) + b_j \otimes 1_k \right\}_{j=m+1}^n$.

Lemma 32. *There exists an isomorphism of S_k -modules (dashed arrow) such that the following diagram is commutative:*

$$\begin{array}{ccc} \mathcal{X}_x \otimes_S S_k & \xrightarrow{\sim} & \mathcal{X}_x(k) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Gamma, k) \end{array}$$

6.3. Description of $\mathcal{X}_x(k)$. We are going to describe this module via the dual of $\mathcal{X}^x(k)$. This is a well known description due to Fiebig [F, Lemmas 6.8, 6.9 and 6.13]. We present here an alternative proof that does not require invertibility of 2 in k . Let

$$D\mathcal{X}^x(k) = \{g \in \text{Map}(\Gamma_x, Q_k) \mid (g, f) \in S_k \text{ for any } f \in \mathcal{X}^x(k)\},$$

where Q_k is the ring of quotients of S_k and $(g, f) = \sum_{\gamma \in \Gamma_x} g_\gamma f_\gamma$ (the standard scalar product). It will be convenient, for example, in Lemma 33 to identify elements of $D\mathcal{X}^x(k)$ with their extensions by zero to Γ .

Consider $P_k \in H_T^\bullet(\Gamma, k)$ defined by

$$P_k(\gamma) = \prod_{i=1}^r \beta_i(\gamma) \otimes 1_k = \pm \prod_{\alpha \in \Phi^+} (\alpha \otimes 1_k)^{|M_\alpha(\gamma)|}.$$

Note that any $P_k(\gamma)$ is divisible in S_k by the Euler class $e_x(k) = \prod_{\alpha \in \Phi^+, s_\alpha x < x} \alpha \otimes 1_k$ (see for example [S, Lemma 4.9.7]).

Lemma 33. $\mathcal{X}_x(k) = P_k D\mathcal{X}^x(k)$.

Proof. First we prove by induction on r that $P_k g \in H_T^\bullet(\Gamma_x, k)$ for any $g \in D\mathcal{X}^x(k)$. This is clear for $r = 0$, so we assume that $r > 0$. Let $t \in \{e, s_r\}$. By Lemma 10, we get

$$S_k \ni (g, \nabla_t f'|_{\Gamma_x}) = \sum_{\gamma' \in \Gamma'_{xt}} g(\gamma' \cdot t) \beta_r(\gamma' \cdot t) f'(\gamma')$$

for any $f' \in \mathcal{X}'(k)$. Hence the function

$$g'(\gamma') = \beta_r(\gamma' \cdot t) g(\gamma' \cdot t)$$

where $\gamma' \in \Gamma'_{xt}$, belongs to $D(\mathcal{X}')^{xt}(k)$. By the inductive hypothesis, the product $P'_k g'$ has values in S_k :

$$S_k \ni P'(\gamma') g'(\gamma') = \left(\prod_{i=1}^{r-1} \beta_i(\gamma') \right) \beta_r(\gamma' \cdot t) g(\gamma' \cdot t) = P(\gamma' \cdot t) g(\gamma' \cdot t).$$

As t is arbitrary, the function $P_k g$ has values in S_k .

Now we are going to prove the lemma $k = \mathbb{Z}'$. In this case, we denote $P = P_{\mathbb{Z}'}$ and $D\mathcal{X}^x = D\mathcal{X}^x(\mathbb{Z}')$. We apply induction on r , the result being obvious for $r = 0$. Assume that $r > 0$.

Let us prove that $Pg \in \mathcal{X}_x$ for $g \in D\mathcal{X}^x$. We actually must prove that the extension by zero of Pg to Γ belongs to $\mathcal{X}$, which by Proposition 7 is equivalent to checking that

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} P(\delta)g(\delta) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}$$

for any $\alpha \in \Phi^+$ and $\gamma \in \Gamma$. In this summation, $P(\delta)$ is clearly divisible by $\alpha^{|M_\alpha(\delta)|} = \alpha^{|M_\alpha(\gamma)|}$. Hence it is also divisible by $\alpha^{|J_\alpha(\gamma)|}$. Therefore it remains to prove that the function

$$p_\gamma^\alpha(\delta) = \begin{cases} (-1)^{|J_\alpha(\delta)|} P(\delta) / \alpha^{|J_\alpha(\gamma)|} & \text{if } \delta \sim_\alpha \gamma \text{ and } J_\alpha(\delta) \subset J_\alpha(\gamma); \\ 0 & \text{otherwise} \end{cases}$$

where $\delta \in \Gamma$, belongs to $\mathcal{X}$. It is rather difficult to prove it directly, applying Proposition 7. Therefore, we define the following function q_γ^α by induction: $q_\gamma^\alpha = 1$ and

$$q_\gamma^\alpha = \begin{cases} \nabla_{\gamma_r} q_{\gamma'}^\alpha & \text{if } r \notin M_\alpha(\gamma); \\ \nabla_{\gamma_r} q_{\gamma'}^\alpha + \nabla_{\gamma_r s_r} q_{\gamma'}^\alpha - \alpha \Delta q_{\gamma'}^\alpha & \text{if } r \in M_\alpha(\gamma) \setminus J_\alpha(\gamma) \\ -\Delta q_{\gamma'}^\alpha & \text{if } r \notin J_\alpha(\gamma). \end{cases}$$

if $r > 0$. By Lemmas 9 and 10, we get $q_\gamma^\alpha \in \mathcal{X}$. Therefore, it suffices to prove that

$$2^{|M_\alpha(\gamma)| - |J_\alpha(\gamma)|} p_\gamma^\alpha = q_\gamma^\alpha.$$

This formula is obvious for $r = 0$. Therefore, we consider the case $r > 0$ and apply induction.

Case 1: $r \notin M_\alpha(\gamma)$. If $\delta \not\sim_\alpha \gamma$, then either $\delta_r \neq \gamma_r$ or $\delta' \not\sim_\alpha \gamma'$. In both cases, $q_\gamma^\alpha(\delta) = \nabla_{\gamma_r} q_{\gamma'}^\alpha(\delta) = 0$. Now assume that $\delta \sim_\alpha \gamma$. Then $\delta_r = \gamma_r$, $\delta' \sim_\alpha \gamma'$, $J_\alpha(\delta) = J_\alpha(\delta')$, $J_\alpha(\gamma) = J_\alpha(\gamma')$, $M_\alpha(\gamma) = M_\alpha(\gamma')$. If $J_\alpha(\delta) \not\subset J_\alpha(\gamma)$, then $J_\alpha(\delta') \not\subset J_\alpha(\gamma')$ and we get

$$q_\gamma^\alpha(\delta) = \nabla_{\gamma_r} q_{\gamma'}^\alpha(\delta) = \beta_r(\delta) q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')|} \beta_r(\delta) p_{\gamma'}^\alpha(\delta') = 0.$$

If $J_\alpha(\delta) \subset J_\alpha(\gamma)$, then $J_\alpha(\delta') \subset J_\alpha(\gamma')$ and we get

$$\begin{aligned} q_\gamma^\alpha(\delta) &= \nabla_{\gamma_r} q_{\gamma'}^\alpha(\delta) = \beta_r(\delta) q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')|} \beta_r(\delta) p_{\gamma'}^\alpha(\delta') \\ &= 2^{|M_\alpha(\gamma)| - |J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta)|} \beta_r(\delta) P(\delta') / \alpha^{|J_\alpha(\gamma)|} \\ &= 2^{|M_\alpha(\gamma)| - |J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta)|} P(\delta) / \alpha^{|J_\alpha(\gamma)|} = 2^{|M_\alpha(\gamma)| - |J_\alpha(\gamma)|} p_\gamma^\alpha(\delta). \end{aligned}$$

Case 2: $r \in M_\alpha(\gamma) \setminus J_\alpha(\gamma)$. If $\delta \not\sim_\alpha \gamma$, then $\delta' \not\sim_\alpha \gamma'$. In this case, $q_\gamma^\alpha(\delta) = 0$, as $q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')|} p_{\gamma'}^\alpha(\delta') = 0$. If $J_\alpha(\delta) \not\subset J_\alpha(\gamma)$, then either $r \in J_\alpha(\delta)$ or $J_\alpha(\delta') \not\subset J_\alpha(\gamma')$. In the former case, we get $\beta_r(\delta) = \alpha$ and

$$q_\gamma^\alpha(\delta) = \nabla_{\gamma_r} q_{\gamma'}^\alpha(\delta) + \nabla_{\gamma_r s_r} q_{\gamma'}^\alpha(\delta) - \alpha \Delta q_{\gamma'}^\alpha(\delta) = \beta_r(\delta) q_{\gamma'}^\alpha(\delta') - \alpha q_{\gamma'}^\alpha(\delta') = 0.$$

In the latter case, we get $q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')|} p_{\gamma'}^\alpha(\delta') = 0$, whence $q_\gamma^\alpha(\delta) = 0$.

Now suppose that $\delta \sim_\alpha \gamma$ and $J_\alpha(\delta) \subset J_\alpha(\gamma)$. Then $\delta' \sim_\alpha \gamma'$, $J_\alpha(\delta') \subset J_\alpha(\gamma')$ and $r \notin J_\alpha(\delta)$. It follows from the last formula that $\beta_r(\delta) = -\alpha$. Hence we get

$$\begin{aligned} q_\gamma^\alpha(\delta) &= \nabla_{\gamma_r} q_{\gamma'}^\alpha(\delta) + \nabla_{\gamma_r s_r} q_{\gamma'}^\alpha(\delta) - \alpha \Delta q_{\gamma'}^\alpha(\delta) = \beta_r(\delta) q_{\gamma'}^\alpha(\delta') - \alpha q_{\gamma'}^\alpha(\delta') \\ &= -2\alpha q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')| + 1} (-1)^{|J_\alpha(\delta')|} (-\alpha) P(\delta') / \alpha^{|J_\alpha(\gamma')|} \\ &= 2^{|M_\alpha(\gamma)| - |J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta)|} P(\delta) / \alpha^{|J_\alpha(\gamma)|} = 2^{|M_\alpha(\gamma)| - |J_\alpha(\gamma)|} p_\gamma^\alpha(\delta). \end{aligned}$$

Case 3: $r \in J_\alpha(\gamma)$. If $\delta \not\sim_\alpha \gamma$, then $\delta' \not\sim_\alpha \gamma'$. In this case, $q_\gamma^\alpha(\delta) = 0$, as $q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')|} p_{\gamma'}^\alpha(\delta') = 0$. If $J_\alpha(\delta) \not\subset J_\alpha(\gamma)$, then $J_\alpha(\delta') \not\subset J_\alpha(\gamma')$ and we again get $q_\gamma^\alpha(\delta) = 0$, as $q_{\gamma'}^\alpha(\delta') = 2^{|M_\alpha(\gamma')| - |J_\alpha(\gamma')|} p_{\gamma'}^\alpha(\delta') = 0$.

Now suppose that $\delta \sim_\alpha \gamma$ and $J_\alpha(\delta) \subset J_\alpha(\gamma)$. Then $\delta' \sim_\alpha \gamma'$ and $J_\alpha(\delta') \subset J_\alpha(\gamma')$.

If $r \notin J_\alpha(\delta)$, then $\beta_r(\delta) = -\alpha$ and

$$\begin{aligned} q_\gamma^\alpha(\delta) &= -\Delta q_\gamma^\alpha(\delta) = -q_\gamma^\alpha(\delta') = -2^{|M_\alpha(\gamma')|-|J_\alpha(\gamma')|} p_\gamma^\alpha(\delta') \\ &= -2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta')|} P(\delta') / \alpha^{|J_\alpha(\gamma')|} = 2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta')|} (-\alpha) P(\delta') / \alpha^{|J_\alpha(\gamma)|} \\ &= 2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta)|} P(\delta) / \alpha^{|J_\alpha(\gamma)|} = 2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} p_\gamma^\alpha(\delta). \end{aligned}$$

If $r \in J_\alpha(\delta)$, then $\beta_r(\delta) = \alpha$ and

$$\begin{aligned} q_\gamma^\alpha(\delta) &= -\Delta q_\gamma^\alpha(\delta) = -q_\gamma^\alpha(\delta') = -2^{|M_\alpha(\gamma')|-|J_\alpha(\gamma')|} p_\gamma^\alpha(\delta') \\ &= -2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta')|} P(\delta') / \alpha^{|J_\alpha(\gamma')|} = 2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta)|} \alpha P(\delta') / \alpha^{|J_\alpha(\gamma)|} \\ &= 2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} (-1)^{|J_\alpha(\delta)|} P(\delta) / \alpha^{|J_\alpha(\gamma)|} = 2^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} p_\gamma^\alpha(\delta). \end{aligned}$$

Finally, we prove the inverse inclusion. Let $f \in \mathcal{X}_x$. We apply induction on the cardinality of the following set (lower closure of the support of f):

$$\widehat{C}(f) = \{\delta \in \Gamma_x \mid \text{there exists } \gamma \in \Gamma_x \text{ such that } \delta \leq \gamma \text{ and } f(\gamma) \neq 0\}.$$

If $\widehat{C}(f) = \emptyset$, then $f = 0$ and the result follows. Suppose now that $\widehat{C}(f) \neq \emptyset$ and let γ be its element maximal with respect to $<$. Let α be a positive root.

First suppose that $s_\alpha x > x$. In this case, $|J_\alpha(\gamma)| = |D_\alpha(\gamma)|$. From [H, Theorem 6.2(3)], we get

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, J_\alpha(\gamma) \subset J_\alpha(\delta)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|}}.$$

Proposition 1 and (21) imply that $\delta \geq \gamma$ for any δ in the above summation. Thus $f(\gamma)$ is divisible by $\alpha^{|M_\alpha(\gamma)|-|J_\alpha(\gamma)|} = \alpha^{|M_\alpha(\gamma)|-|D_\alpha(\gamma)|}$.

Now suppose that $s_\alpha x < x$. In this case, $|J_\alpha(\gamma)| = |D_\alpha(\gamma)| + 1$. Let j be the greatest element of $M_\alpha(\gamma)$. Note that j is also the greatest element of $J_\alpha(\gamma)$. Let $\tilde{\gamma}$ be obtained from γ by replacing γ_j with $\gamma_j s_j$. We clearly have $\pi(\tilde{\gamma}) = s_\alpha x$, $\tilde{\gamma} \sim_\alpha \gamma$ and $J_\alpha(\gamma) = J_\alpha(\tilde{\gamma}) \sqcup \{j\}$, whence $|J_\alpha(\tilde{\gamma})| = |J_\alpha(\gamma)| - 1 = |D_\alpha(\gamma)|$. From [H, Theorem 6.2(3)], we get

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, J_\alpha(\tilde{\gamma}) \subset J_\alpha(\delta)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|M_\alpha(\gamma)|-|D_\alpha(\gamma)|}}.$$

As $j \in J_\alpha(\delta)$ for any δ in the above summation, we can replace there the condition $J_\alpha(\tilde{\gamma}) \subset J_\alpha(\delta)$ with $J_\alpha(\gamma) \subset J_\alpha(\delta)$. Hence again $\delta \geq \gamma$ in the above summation and $f(\gamma)$ is divisible by $\alpha^{|M_\alpha(\gamma)|-|D_\alpha(\gamma)|}$.

As a result, we get that $f(\gamma)$ is divisible by

$$\prod_{\alpha \in \Phi^+} \alpha^{|M_\alpha(\gamma)|-|D_\alpha(\gamma)|} = \pm \frac{P(\gamma)}{\mathbf{a}(\gamma)}.$$

It follows from Theorem 17 that $D\mathcal{X}^x$ has an S -basis $\{\hat{\mathbf{b}}_\gamma\}_{\gamma \in \Gamma_x}$ such that $\hat{\mathbf{b}}_\gamma(\gamma) = 1/\mathbf{a}(\gamma)$ and $\hat{\mathbf{b}}_\gamma(\delta) = 0$ for $\delta \in \Gamma_x$ with $\delta > \gamma$. To get this basis, one should invert and transpose the matrix given by (22).

Consider the difference $h = f - f(\gamma)/(P(\gamma)/\mathbf{a}(\gamma))P\hat{\mathbf{b}}_\gamma$. We get $C(h) \subset \{\delta \in \Gamma_x \mid \delta < \gamma\} \subsetneq C(f)$. By induction, h belongs to the $PD\mathcal{X}^x$. Thus so does f .

Finally, let us return to the general case. By Lemma 28, the basis $\{\hat{\mathbf{b}}_\gamma\}_{\gamma \in \Gamma_x}$ of $D\mathcal{X}^x$ mentioned above and the similar basis for $D\mathcal{X}^x(k)$ yield an isomorphism (dashed arrow)

making the following diagram commutative:

$$\begin{array}{ccc} PD\mathcal{X}^x \otimes_S S_k & \xrightarrow{\sim} & P_k D\mathcal{X}^x(k) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Gamma, k) \end{array}$$

Multiplying it by P_k , we get by Lemma 32, an isomorphism (dashed arrow) making the following diagram commutative:

$$\begin{array}{ccc} & \mathcal{X}_x \otimes_S S_k & \xrightarrow{\sim} \mathcal{X}_x(k) \\ & \nearrow & \nearrow \\ (PD\mathcal{X}^x) \otimes_S S_k & \xrightarrow{\sim} & P_k D\mathcal{X}^x(k) \\ \downarrow & & \downarrow \\ H_T^\bullet(\Gamma) \otimes_S S_k & \xrightarrow{\sim} & H_T^\bullet(\Gamma, k) \end{array}$$

(Note: A dashed arrow also connects $\mathcal{X}_x(k)$ to $H_T^\bullet(\Gamma, k)$ in the original diagram.)

The commutativity of the right triangle means that the isomorphism represented by the dashed arrow is over $H_T^\bullet(\Gamma, k)$, which proves that it is the equality of subsets. $\square$

Proposition 34. *Let H_x be the matrix defined by (22) and P_x be the diagonal matrix with γ^{th} entry $P_{\mathbb{Z}'}(\gamma)$. We set $H_{x,k} = H_x \otimes_S S_k$ and $P_{x,k} = P_x \otimes_S S_k$. The costalk-to-stalk embedding $\mathcal{X}_x(k) \hookrightarrow \mathcal{X}^x(k)$ is described by the transition matrix $(H_{x,k}^{-1})^T P_{x,k} H_{x,k}^{-1}$. All entries of this matrix are divisible in S_k by the Euler class $e_x(k) = \prod_{\alpha \in \Phi^+, s_\alpha x < x} \alpha \otimes 1_k$.*

Proof. We need only to prove the divisibility. It suffices to consider the case $k = \mathbb{Z}'$. We denote $e_x = e_x(\mathbb{Z}')$. Let $f \in \mathcal{X}_x$. By Proposition 7, we get

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, J_\alpha(\delta) \subset J_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\delta) \equiv 0 \pmod{\alpha^{|J_\alpha(\gamma)|}}$$

for any $\alpha \in \Phi^+$. If $s_\alpha x < x$, then $|J_\alpha(\gamma)| = |D_\alpha(\gamma)| + 1$. By Proposition 1 and (21), we get $\delta \leq \gamma$ in the above summation. Hence we get by induction that $f(\gamma)$ is divisible in S by $e_x = e_x(\mathbb{Z}')$. Dividing the above equivalence by α if $s_\alpha x < x$ and taking into account that different roots are not proportional, we obtain

$$\sum_{\delta \in \Gamma_x, \delta \sim_\alpha \gamma, D_\alpha(\delta) \subset D_\alpha(\gamma)} (-1)^{|J_\alpha(\delta)|} f(\delta)/e_x \equiv 0 \pmod{\alpha^{|D_\alpha(\gamma)|}}.$$

Thus we have proved that $f/e_x \in \mathcal{X}^x$. $\square$

6.4. Euler classes. Consider the following closed T -equivariant embedding

$$f : \{0\} \hookrightarrow V = \mathbb{C}_{\lambda_1} \oplus \cdots \oplus \mathbb{C}_{\lambda_d},$$

where λ_i are characters of T and $\mathbb{C}_{\lambda_i}$ is the corresponding one-dimensional representation of T (see the proof of Theorem 5). We assume that k is a field such that $\lambda_i \otimes_{\mathbb{Z}} k \neq 0$ for any i . We get the following exact sequence

$$H_T^\bullet(\{0\}, f^! \underline{k}_V) \rightarrow H_T^\bullet(V, k) \rightarrow H_T^\bullet(V \setminus \{0\}, k) \quad (43)$$

We want to look more closely at the right map. By [AB, (2.15)], we have the following commutative diagram with exact top row:

$$\begin{array}{ccccc} H_T^\bullet(V, V \setminus \{0\}, k) & \longrightarrow & H_T^\bullet(V, k) & \longrightarrow & H_T^\bullet(V \setminus \{0\}, k) \\ \uparrow \wr \Phi & \nearrow f_* & \downarrow \wr f^* & & \\ H_T^{\bullet-2d}(\{0\}, k) & \xrightarrow{e \cup ?} & H_T^\bullet(\{0\}, k) & & \end{array}$$

where Φ is the Thom isomorphism, f_* is the push-forward and $e = \prod_{i=1}^d \lambda_i \otimes_{\mathbb{Z}} k$ is the corresponding Euler class. By our assumption $e \neq 0$. Hence $H_T^\bullet(V \setminus \{0\}, k)$ vanishes in odd degrees and $H_T^\bullet(V, k) \rightarrow H_T^\bullet(V \setminus \{0\}, k)$ is epimorphic in any degree. Coming back to (43), we obtain the isomorphisms $H_T^\bullet(\{0\}, f^! \underline{k}_V) \simeq H_T^{\bullet-2d}(\{0\}, k) \simeq S_k[-2d]$ and $H_T^\bullet(V, k) \simeq S_k$ under which the map $H_T^\bullet(\{0\}, f^! \underline{k}_V) \rightarrow H_T^\bullet(V, k)$ becomes the multiplication by e .

6.5. Defect of a homomorphism. Recall that S_k has the maximal ideal $\mathfrak{m} = \bigoplus_{i>0} S_k^i$. We clearly have $S_k/\mathfrak{m} \simeq k$. Let $\varphi : U \rightarrow V$ be a homomorphism of graded S_k -modules. Then we can consider the quotient $\text{im } \varphi / \mathfrak{m}V$, which is a graded $S_k/\mathfrak{m}$ -module and thus also a graded k -vector space. If U is a finitely generated S_k -module, then we can define the *defect* of φ as the graded dimension of this quotient:

$$d(\varphi) = \sum_{n \in \mathbb{Z}} \dim_k(\text{im } \varphi / \mathfrak{m}V)^n v^{-n}.$$

This is an element of the ring of Laurent polynomials $\mathbb{Z}[v, v^{-1}]$. Clearly $d(\varphi_1 \oplus \varphi_2) = d(\varphi_1) + d(\varphi_2)$ and $d(\varphi[n]) = v^n d(\varphi)$. If φ is an embedding and U and V are finitely generated free S_k -modules, then $d(\varphi)$ can be calculated as follows.

Proposition 35 ([S, Corollary 3.3.3]). *Let $\varphi : U \hookrightarrow V$ be an embedding of graded S_k -modules. Suppose that $\{u_i^{(n)}\}_{n \in \mathbb{Z}, 1 \leq i \leq l_n}$ and $\{v_j^{(n)}\}_{n \in \mathbb{Z}, 1 \leq j \leq k_n}$ are bases of U and V respectively labelled in such a way that $u_i^{(n)}$ and $v_j^{(n)}$ have degree n . Let*

$$\varphi(u_i^{(n)}) = \sum_{m \in \mathbb{Z}, 1 \leq j \leq k_m} a_{j,i}^{(m,n)} v_j^{(m)}$$

for corresponding homogeneous $a_{j,i}^{(m,n)} \in S_k$. For each $n \in \mathbb{Z}$, we denote by $A^{(n)}$ the $k_n \times l_n$ -matrix whose ji^{th} entry is $a_{j,i}^{(n,n)} \in k$. Then $d(\varphi) = \sum_{n \in \mathbb{Z}} \text{rk}_k A^{(n)} v^{-n}$.

Finally, we describe a homomorphism of graded modules $\varphi : U \rightarrow V$ can be divided by a homogeneous element $e \in S_k^d$ that is no zero divisor for V . Suppose that for any $u \in U$ there exists $(\varphi/e)(u) \in V$ such that $\varphi(u) = e(\varphi/e)(u)$. Then we get a uniquely defined homomorphism $\varphi/e : U \rightarrow V$ of S_k -modules such that $(\varphi/e)(U_{i+d}) \subset V_i$. It is a homomorphism of graded modules $\varphi/e : U[d] \rightarrow V$.

6.6. Application to parity sheaves. Our calculations have not yet involved any stratifications. In this section, we are going to apply our results to the stratification $G/B = \bigsqcup_{x \in W} BxB/B$. We denote $X = G/B$ and $X_x = BxB/B$ for brevity and assume that k is a field. By [JMW] there exists the following decomposition:

$$\pi_* \underline{k}_\Sigma[r] = \bigoplus_{x \in W} \bigoplus_{d \in \mathbb{Z}} \mathcal{E}(x, k)[-d]^{\oplus m(x,d)},$$

where $\mathcal{E}(x, k) \in D_T(X, k)$ is the T -equivariant parity sheaf such that $\text{supp } \mathcal{E}(x, k) \subset \overline{X_x}$ and $\mathcal{E}(x, k)|_{X_x} = \underline{k}_{X_x}[d_x]$, where $d_x = \dim X_x$. Our aim is to calculate the multiplicities $m(x, d)$.

We rewrite the above decomposition as follows:

$$\pi_* \underline{k}_\Sigma = \bigoplus_{x \in W} \bigoplus_{d \in \mathbb{Z}} \mathcal{E}(x, k)[-d-r]^{\oplus m(x,d)} \quad (44)$$

and consider the natural embedding $i_x : \{x\} \hookrightarrow X$. We are going to take the following steps:

- apply to both sides of (44) the morphism of functors $H_T^\bullet(\{x\}, i_x^! -) \rightarrow H_T^\bullet(\{x\}, i_x^* -)$;
- divide it by the Euler class $e_x = \prod_{\alpha \in \Phi^+, s_\alpha x < x} \alpha \otimes 1_k$;
- take the defect of the resulting map.

First consider the left-hand side of (44). We have the following Cartesian diagram:

$$\begin{array}{ccc} \Sigma_x & \xrightarrow{i} & \Sigma \\ \downarrow \pi_x & & \downarrow \pi \\ \{x\} & \xrightarrow{i_x} & X \end{array}$$

As π is proper, the base change yields $i_x^! \pi_* \underline{k}_\Sigma \simeq (\pi_x)_* i_x^! \underline{k}_\Sigma$ and $i_x^* \pi_* \underline{k}_\Sigma \simeq (\pi_x)_* i_x^* \underline{k}_\Sigma$. Hence the map $H_T^\bullet(\{x\}, i_x^! \pi_* \underline{k}_\Sigma) \rightarrow H_T^\bullet(\{x\}, i_x^* \pi_* \underline{k}_\Sigma)$ is isomorphic to $H_T^\bullet(\Sigma_x, i_x^! \underline{k}_\Sigma) \rightarrow H_T^\bullet(\Sigma_x, i_x^* \underline{k}_\Sigma)$, which in its turn is isomorphic to $\mathcal{X}_x(k) \hookrightarrow \mathcal{X}^x(k)$ by Corollary 31.

In order to tackle the right-hand side of (44), let us compute the map

$$H_T^\bullet(\{x\}, i_x^! \mathcal{E}(y, k)) \rightarrow H_T^\bullet(\{x\}, i_x^* \mathcal{E}(y, k)). \quad (45)$$

If $x \notin \overline{X_y}$, then $i_x^! \mathcal{E}(y, k) = i_x^* \mathcal{E}(y, k) = 0$ as $\mathcal{E}(y, k)|_{X \setminus \overline{X_y}} = 0$. Therefore, we must only consider the case $x \in \overline{X_y}$ that is $x \leq y$.

Let $U = \bigsqcup_{z \geq x} X_z$. This is an open subset of X , which contains X_x as a closed subset. Moreover, the restriction $\mathcal{F} = \mathcal{E}(y, k)|_U$ is an indecomposable parity sheaf on U . Let $i : X_x \hookrightarrow U$, $f : \{x\} \hookrightarrow X_x$ and $\tilde{i}_x : \{x\} \hookrightarrow U$ denote the natural embeddings:

$$\begin{array}{ccccc} & & \tilde{i}_x & & \\ & \nearrow & & \searrow & \\ \{x\} & \xrightarrow{f} & X_x & \xrightarrow{i} & U \longrightarrow X \\ & \searrow & & \nearrow & \\ & & i_x & & \end{array}$$

Let $\varphi : i^! \mathcal{F} \rightarrow i^* \mathcal{F}$ denote the natural morphism.

Suppose first that $x < y$. Then $\mathcal{F}$ has no direct summands supported on X_x . Let us write $i^! \mathcal{F} = \bigoplus_{m \in \mathbb{Z}} Q^m$ and $i^* \mathcal{F} = \bigoplus_{n \in \mathbb{Z}} P^n$, where $Q^m = H^m(i^! \mathcal{F})[-m]$ and $P^n = H^n(i^* \mathcal{F})[-n]$. We denote by $\varphi_{n,m} : Q^m \rightarrow P^n$ the corresponding morphism of the direct summands. By [JMW, Corollary 2.22], we get that $\varphi_{n,m} = 0$ for $m \leq n$.

We denote by $A : f^! \rightarrow f^*$ the natural morphism of functors. Then the composition $f^* \varphi \circ A(i^! \mathcal{F})$ is the natural morphism $\tilde{i}_x^! \mathcal{F} \rightarrow \tilde{i}_x^* \mathcal{F}$:

$$\tilde{i}_x^! \mathcal{F} = f^! \circ i^! \mathcal{F} \xrightarrow{A(i^! \mathcal{F})} f^* \circ i^! \mathcal{F} \xrightarrow{f^* \varphi} f^* \circ i^* \mathcal{F} = \tilde{i}_x^* \mathcal{F}.$$

Recalling our decompositions of $i^! \mathcal{F}$ and $i^* \mathcal{F}$, we represent this morphism as the direct sum of the following morphisms:

$$f^! Q^m \xrightarrow{A(Q^m)} f^* Q^m \xrightarrow{f^* \varphi_{n,m}} f^* P^n \quad (46)$$

for $m > n$. We have the decompositions $Q^m = \underline{k}_{X_x}[-m]^{\oplus a(x,m)}$ and $P^n = \underline{k}_{X_x}[-n]^{\oplus b(x,n)}$ for some nonnegative integers $a(x,m)$ and $b(x,n)$. Applying $H_T^\bullet(\{x\}, -)$ to (46), we get maps from

$$\begin{aligned} H_T^\bullet(\{x\}, f^! Q^m) &= H_T^\bullet(\{x\}, f^! \underline{k}_{X_x}[-m])^{\oplus a(x,m)} = H_T^{\bullet-m}(\{x\}, f^! \underline{k}_{X_x})^{\oplus a(x,m)} \\ &= H_T^{\bullet-m-2d_x}(\{x\}, k)^{\oplus a(x,m)} = S_k[-m-2d_x]^{\oplus a(x,m)} \end{aligned}$$

to

$$H_T^\bullet(\{x\}, f^* Q^m) = H_T^\bullet(\{x\}, f^* \underline{k}_{X_x}[-m])^{\oplus a(x,m)} = H_T^{\bullet-m}(\{x\}, k)^{\oplus a(x,m)} = S_k[-m]^{\oplus a(x,m)}$$

and finally to

$$H_T^\bullet(\{x\}, f^* P^n) = H_T^\bullet(\{x\}, f^* \underline{k}_{X_x}[-n])^{\oplus b(x,n)} = H_T^{\bullet-n}(\{x\}, k)^{\oplus b(x,n)} = S_k[-n]^{\oplus b(x,n)}.$$

So we get the following sequence of maps (see Section 6.4):

$$S_k[-m-2d_x]^{\oplus a(x,m)} \xrightarrow{e_x \cup ?} S_k[-m]^{\oplus a(x,m)} \longrightarrow S_k[-n]^{\oplus b(x,n)}.$$

The division by e_x yields a map $S_k[-m]^{\oplus a(x,m)} \rightarrow S_k[-n]^{\oplus b(x,n)}$. As $m > n$ the defect of this map is zero. Thus we have proved that the defect of the map $H_T^\bullet(\{x\}, \tilde{i}_x^! \mathcal{F}) \rightarrow H_T^\bullet(\{x\}, \tilde{i}_x^* \mathcal{F})$ divided by e_x is zero. Recalling that $\mathcal{F} = \mathcal{E}(y, k)|_U$, we get that the defect of (45) divided by e_x is also zero.

It remains to calculate the defect of (45) divided by e_x in the case $x = y$. Consider the natural embedding $j : U \setminus X_x \hookrightarrow U$. We have $j^* \mathcal{F} = \mathcal{E}(x, k)|_{U \setminus X_x} = 0$ as $U \setminus X_x \subset X \setminus \overline{X_x}$. The distinguished triangle

$$0 = j_! j^* \mathcal{F} \rightarrow \mathcal{F} \rightarrow i_* i^* \mathcal{F} \xrightarrow{+1}$$

yields $\mathcal{F} = i_* i^* \mathcal{F} = i_* \underline{k}_{X_x}[d_x]$. For any $? \in \{!, *\}$, we get

$$i_x^? \mathcal{E}(x, k) = \tilde{i}_x^? \mathcal{F} = \tilde{i}_x^? i_* \underline{k}_{X_x}[d_x] = f^? i^? i_* \underline{k}_{X_x}[d_x] = f^? \underline{k}_{X_x}[d_x].$$

Hence (45) becomes the natural map

$$H_T^{\bullet+d_x}(\{x\}, f^! \underline{k}_{X_x}) \rightarrow H_T^{\bullet+d_x}(\{x\}, f^* \underline{k}_{X_x}).$$

Under the identifications of Section 6.4, this map becomes $S[-d_x] \xrightarrow{e_x \cup ?} S[d_x]$. Division by e_x leaves us with the identity map $S_k[d_x] \rightarrow S_k[d_x]$, whose defect is obviously is v^{d_x} . Hence we get the following result.

Theorem 36. *The defect of the inclusion $\mathcal{X}_x(k) \hookrightarrow \mathcal{X}^x(k)$ divided by e_x is*

$$\sum_{d \in \mathbb{Z}} m(x, d) v^{d_x - d - r}.$$

Once we compute the above inclusion, for example, by Proposition 34, we can recover the coefficients $m(x, d)$.

6.7. Example of torsion. We use here the notation of Proposition 34. Let $G = \mathrm{SL}_8(\mathbb{C})$, $\Pi = \{\alpha_1, \dots, \alpha_7\}$ and

$$s = (s_3, s_2, s_1, s_5, s_4, s_3, s_2, s_6, s_5, s_4, s_3, s_7, s_6, s_5), \quad x = s_2 s_3 s_2 s_5 s_6 s_5,$$

where $s_i = s_{\alpha_i}$. We arrange elements of Γ_x in ascending order with respect to $<$. The matrix $H_x = \{h_{i,j}\}_{i,j=1}^{29}$ computed by (22) has the following nonzero entries:

$$\begin{aligned}
& h_{1,j} = 1 \text{ for } 1 \leq j \leq 29, h_{13,13} = h_{13,14} = h_{13,15} = h_{13,16} = h_{13,17} = h_{13,18} = h_{13,19} = \alpha_5 + \alpha_6, \\
& h_{3,3} = h_{3,6} = h_{3,8} = h_{3,15} = h_{3,17} = h_{3,22} = h_{3,27} = \alpha_2 + \alpha_3, h_{7,7} = h_{7,8} = h_{7,16} = h_{7,17} = \alpha_3 + \alpha_4 + \alpha_5, \\
& h_{20,20} = h_{20,21} = h_{20,22} = h_{20,23} = h_{20,24} = h_{20,25} = h_{20,26} = h_{20,27} = h_{20,28} = h_{20,29} = \alpha_6, \\
& h_{4,4} = h_{4,5} = h_{4,6} = h_{4,11} = h_{4,12} = h_{13,25} = h_{13,26} = h_{13,27} = h_{13,28} = h_{13,29} = \alpha_5, h_{27,27} = \alpha_6 \alpha_5 (\alpha_2 + \alpha_3) \\
& h_{9,9} = h_{9,10} = h_{9,11} = h_{9,12} = h_{9,18} = h_{9,19} = h_{9,23} = h_{9,24} = h_{9,28} = h_{9,29} = \alpha_2, h_{5,14} = h_{6,19} = -\alpha_3 \alpha_6, \\
& h_{2,2} = h_{2,5} = h_{2,14} = h_{2,21} = h_{2,26} = h_{3,10} = h_{3,12} = h_{3,19} = h_{3,24} = h_{3,29} = \alpha_3, h_{26,27} = -\alpha_6 \alpha_2 \alpha_5, \\
& h_{4,16} = h_{4,17} = -\alpha_3 - \alpha_4 - \alpha_5 - \alpha_6, h_{2,7} = h_{2,16} = -\alpha_4 - \alpha_5, h_{2,8} = h_{2,17} = -\alpha_2 - \alpha_3 - \alpha_4 - \alpha_5, \\
& h_{12,12} = h_{19,29} = \alpha_2 \alpha_3 \alpha_5, h_{26,26} = h_{27,29} = \alpha_6 \alpha_3 \alpha_5, h_{6,17} = -(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_2 + \alpha_3), \\
& h_{5,6} = h_{14,27} = -\alpha_2 \alpha_5, h_{10,10} = h_{10,12} = h_{10,19} = h_{10,24} = h_{10,29} = \alpha_2 \alpha_3, h_{28,28} = h_{28,29} = \alpha_6 \alpha_2 \alpha_5, \\
& h_{24,24} = h_{24,29} = \alpha_2 \alpha_3 \alpha_6, h_{5,5} = h_{6,12} = h_{14,26} = h_{15,29} = \alpha_3 \alpha_5, h_{11,18} = h_{11,19} = h_{21,22} = h_{21,27} = -\alpha_2 \alpha_6, \\
& h_{22,22} = h_{22,27} = \alpha_6(\alpha_2 + \alpha_3), h_{14,14} = h_{15,19} = (\alpha_5 + \alpha_6) \alpha_3, h_{2,3} = h_{2,6} = h_{2,15} = h_{2,22} = h_{2,27} = -\alpha_2, \\
& h_{6,15} = -\alpha_6(\alpha_2 + \alpha_3), h_{19,19} = (\alpha_5 + \alpha_6) \alpha_2 \alpha_3, h_{15,15} = h_{15,17} = (\alpha_5 + \alpha_6)(\alpha_2 + \alpha_3), h_{14,15} = -(\alpha_5 + \alpha_6) \alpha_2, \\
& h_{14,16} = -(\alpha_5 + \alpha_6)(\alpha_4 + \alpha_5), h_{5,7} = (\alpha_3 + \alpha_4)(\alpha_4 + \alpha_5), h_{14,17} = -(\alpha_5 + \alpha_6)(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5), \\
& h_{18,18} = h_{18,19} = (\alpha_5 + \alpha_6) \alpha_2, h_{5,8} = (\alpha_3 + \alpha_4)(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5), h_{11,11} = h_{11,12} = h_{18,28} = h_{18,29} = \alpha_2 \alpha_5, \\
& h_{16,16} = h_{16,17} = (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4 + \alpha_5), h_{12,19} = -\alpha_2 \alpha_3 \alpha_6, h_{29,29} = \alpha_2 \alpha_3 \alpha_5 \alpha_6, h_{4,7} = h_{4,8} = -\alpha_3 - \alpha_4, \\
& h_{4,13} = h_{4,14} = h_{4,15} = h_{4,18} = h_{4,19} = -\alpha_6, h_{5,15} = h_{23,23} = h_{23,24} = h_{23,28} = h_{23,29} = \alpha_2 \alpha_6, \\
& h_{6,6} = h_{15,27} = \alpha_5(\alpha_2 + \alpha_3), h_{8,8} = h_{8,17} = (\alpha_3 + \alpha_4 + \alpha_5)(\alpha_2 + \alpha_3), h_{17,17} = (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4 + \alpha_5)(\alpha_2 + \alpha_3), \\
& h_{6,8} = -(\alpha_3 + \alpha_4)(\alpha_2 + \alpha_3), h_{5,16} = (\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_4 + \alpha_5), h_{5,17} = (\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5), \\
& h_{21,21} = h_{21,26} = h_{22,24} = h_{22,29} = \alpha_3 \alpha_6, h_{25,25} = h_{25,26} = h_{25,27} = h_{25,28} = h_{25,29} = \alpha_5 \alpha_6.
\end{aligned}$$

The 29th row of the matrix $(H_x^{-1})^T$ has minimal degree. Its precise value is the following:

$$\begin{aligned}
r_{29} = & \left(\frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, -\frac{1}{(\alpha_2 + \alpha_3) \alpha_6 \alpha_5 \alpha_3}, -\frac{1}{\alpha_2 \alpha_6 \alpha_5 (\alpha_2 + \alpha_3)}, -\frac{1}{(\alpha_5 + \alpha_6) \alpha_2 \alpha_5 \alpha_3}, \frac{1}{(\alpha_5 + \alpha_6) (\alpha_2 + \alpha_3) \alpha_5 \alpha_3}, \right. \\
& \frac{1}{\alpha_2 (\alpha_5 + \alpha_6) \alpha_5 (\alpha_2 + \alpha_3)}, 0, 0, -\frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, \frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, \frac{1}{(\alpha_5 + \alpha_6) \alpha_2 \alpha_5 \alpha_3}, -\frac{1}{(\alpha_5 + \alpha_6) \alpha_2 \alpha_5 \alpha_3}, -\frac{1}{\alpha_6 (\alpha_5 + \alpha_6) \alpha_2 \alpha_3}, \\
& \frac{1}{\alpha_6 (\alpha_2 + \alpha_3) (\alpha_5 + \alpha_6) \alpha_3}, \frac{1}{\alpha_2 \alpha_6 (\alpha_5 + \alpha_6) (\alpha_2 + \alpha_3)}, 0, 0, \frac{1}{\alpha_6 (\alpha_5 + \alpha_6) \alpha_2 \alpha_3}, -\frac{1}{\alpha_6 (\alpha_5 + \alpha_6) \alpha_2 \alpha_3}, -\frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, \\
& \frac{1}{(\alpha_2 + \alpha_3) \alpha_6 \alpha_5 \alpha_3}, \frac{1}{\alpha_2 \alpha_6 \alpha_5 (\alpha_2 + \alpha_3)}, \frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, -\frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, \frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, -\frac{1}{(\alpha_2 + \alpha_3) \alpha_6 \alpha_5 \alpha_3}, -\frac{1}{\alpha_2 \alpha_6 \alpha_5 (\alpha_2 + \alpha_3)}, \\
& \left. -\frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3}, \frac{1}{\alpha_6 \alpha_2 \alpha_5 \alpha_3} \right).
\end{aligned}$$

We have the following Euler class: $e_x = \alpha_3(\alpha_2 + \alpha_3) \alpha_6 \alpha_2 (\alpha_5 + \alpha_6) \alpha_5$. Hence the matrix P_x/e_x has the following diagonal:

$$\begin{aligned}
p := & (\alpha_3 \alpha_6 \alpha_2 \alpha_5 \alpha_1 \alpha_4 (\alpha_3 + \alpha_4 + \alpha_5) \alpha_7, -\alpha_3 (\alpha_2 + \alpha_3) \alpha_1 \alpha_5 (\alpha_3 + \alpha_4) \alpha_6 (\alpha_3 + \alpha_4 + \alpha_5) \alpha_7, \\
& -(\alpha_2 + \alpha_3) (\alpha_1 + \alpha_2 + \alpha_3) \alpha_5 (\alpha_3 + \alpha_4) \alpha_2 \alpha_6 (\alpha_3 + \alpha_4 + \alpha_5) \alpha_7, -\alpha_3 \alpha_2 \alpha_1 \alpha_5 (\alpha_4 + \alpha_5) (\alpha_5 + \alpha_6) (\alpha_3 + \alpha_4 + \alpha_5) \alpha_7, \\
& \alpha_3 (\alpha_2 + \alpha_3) \alpha_1 \alpha_5 (\alpha_3 + \alpha_4 + \alpha_5)^2 (\alpha_5 + \alpha_6) \alpha_7, (\alpha_2 + \alpha_3) (\alpha_1 + \alpha_2 + \alpha_3) \alpha_5 (\alpha_3 + \alpha_4 + \alpha_5)^2 \alpha_2 (\alpha_5 + \alpha_6) \alpha_7, \\
& -(\alpha_2 + \alpha_3) \alpha_1 (\alpha_3 + \alpha_4 + \alpha_5)^2 (\alpha_4 + \alpha_5) (\alpha_5 + \alpha_6) (\alpha_3 + \alpha_4) \alpha_7, \\
& (\alpha_2 + \alpha_3) (\alpha_1 + \alpha_2 + \alpha_3) (\alpha_3 + \alpha_4 + \alpha_5)^2 (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5) (\alpha_5 + \alpha_6) (\alpha_3 + \alpha_4) \alpha_7, \\
& -\alpha_3 \alpha_2 (\alpha_1 + \alpha_2) \alpha_5 \alpha_4 \alpha_6 (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5) \alpha_7, \alpha_3 (\alpha_1 + \alpha_2 + \alpha_3) \alpha_5 (\alpha_3 + \alpha_4) \alpha_2 \alpha_6 (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5) \alpha_7, \\
& \alpha_3 \alpha_2 (\alpha_1 + \alpha_2) \alpha_5 (\alpha_4 + \alpha_5) (\alpha_5 + \alpha_6) (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5) \alpha_7, \\
& -\alpha_3 (\alpha_1 + \alpha_2 + \alpha_3) \alpha_5 (\alpha_3 + \alpha_4 + \alpha_5) \alpha_2 (\alpha_5 + \alpha_6) (\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5) \alpha_7, \\
& -\alpha_3 \alpha_2 \alpha_1 (\alpha_4 + \alpha_5) (\alpha_5 + \alpha_6) \alpha_6 (\alpha_3 + \alpha_4 + \alpha_5) (\alpha_5 + \alpha_6 + \alpha_7), \alpha_3 (\alpha_2 + \alpha_3) \alpha_1 (\alpha_3 + \alpha_4 + \alpha_5)^2 (\alpha_5 + \alpha_6) \alpha_6 (\alpha_5 + \alpha_6 + \alpha_7), \\
& (\alpha_2 + \alpha_3) (\alpha_1 + \alpha_2 + \alpha_3) (\alpha_3 + \alpha_4 + \alpha_5)^2 \alpha_2 (\alpha_5 + \alpha_6) \alpha_6 (\alpha_5 + \alpha_6 + \alpha_7),
\end{aligned}$$

$$\begin{aligned}
& (\alpha_2 + \alpha_3)\alpha_1(\alpha_3 + \alpha_4 + \alpha_5)^2(\alpha_4 + \alpha_5)(\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7), \\
& - (\alpha_2 + \alpha_3)(\alpha_1 + \alpha_2 + \alpha_3)(\alpha_3 + \alpha_4 + \alpha_5)^2(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5)(\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7), \\
& \quad \alpha_3\alpha_2(\alpha_1 + \alpha_2)(\alpha_4 + \alpha_5)(\alpha_5 + \alpha_6)\alpha_6(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5)(\alpha_5 + \alpha_6 + \alpha_7), \\
& \quad - \alpha_3(\alpha_1 + \alpha_2 + \alpha_3)(\alpha_3 + \alpha_4 + \alpha_5)\alpha_2(\alpha_5 + \alpha_6)\alpha_6(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5)(\alpha_5 + \alpha_6 + \alpha_7), \\
& - \alpha_3\alpha_2\alpha_1\alpha_5\alpha_4\alpha_6(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_6 + \alpha_7), \alpha_3(\alpha_2 + \alpha_3)\alpha_1\alpha_5(\alpha_3 + \alpha_4)\alpha_6(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_6 + \alpha_7), \\
& \quad (\alpha_2 + \alpha_3)(\alpha_1 + \alpha_2 + \alpha_3)\alpha_5(\alpha_3 + \alpha_4)\alpha_2\alpha_6(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_6 + \alpha_7), \\
& \quad \alpha_3\alpha_2(\alpha_1 + \alpha_2)\alpha_5\alpha_4\alpha_6(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_6 + \alpha_7), \\
& \quad - \alpha_3(\alpha_1 + \alpha_2 + \alpha_3)\alpha_5(\alpha_3 + \alpha_4)\alpha_2\alpha_6(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_6 + \alpha_7), \\
& \quad \alpha_3\alpha_2\alpha_1\alpha_5(\alpha_4 + \alpha_5)\alpha_6(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7), \\
& \quad - \alpha_3(\alpha_2 + \alpha_3)\alpha_1\alpha_5(\alpha_3 + \alpha_4 + \alpha_5)\alpha_6(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7), \\
& \quad - (\alpha_2 + \alpha_3)(\alpha_1 + \alpha_2 + \alpha_3)\alpha_5(\alpha_3 + \alpha_4 + \alpha_5)\alpha_2\alpha_6(\alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7), \\
& \quad - \alpha_3\alpha_2(\alpha_1 + \alpha_2)\alpha_5(\alpha_4 + \alpha_5)\alpha_6(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7), \\
& \quad \alpha_3(\alpha_1 + \alpha_2 + \alpha_3)\alpha_5(\alpha_3 + \alpha_4 + \alpha_5)\alpha_2\alpha_6(\alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_6)(\alpha_5 + \alpha_6 + \alpha_7)
\end{aligned}$$

and zeros elsewhere.

Consider the triple scalar product of rows: $(a, b, c) = \sum_{i=1}^{29} a_i b_i c_i$. A (computer) calculation shows that $(r_{29}, r_{29}, p) = 2$. Thus we have proved that the defect of the inclusion $\mathcal{X}_x(k) \hookrightarrow \mathcal{X}^x(k)$ divided by $e_x(k)$ is 0 if $\text{char } k = 2$ and is v^{-8} otherwise. By Theorem 36, we get

$$\sum_{d \in \mathbb{Z}} m(x, d) v^{-d-8} = \begin{cases} 0 & \text{if } \text{char } k = 2; \\ v^{-8} & \text{otherwise.} \end{cases}$$

Hence we get the following result.

Theorem 37. *Let $G = \text{SL}_8(\mathbb{C})$ and $\Pi = \{\alpha_1, \dots, \alpha_7\}$ be the set of simple roots. Consider the Bott-Samelson variety Σ for the sequence $s = (s_3, s_2, s_1, s_5, s_4, s_3, s_2, s_6, s_5, s_4, s_3, s_7, s_6, s_5)$ and take $x = s_2 s_3 s_2 s_5 s_6 s_5$. Let $\Sigma \rightarrow G/B$ be the canonical resolution and k be a field. If $\text{char } k = 2$, then $\pi_* \underline{k}_\Sigma[14]$ has no direct summand of the form $\mathcal{E}(x, k)[d]$. If $\text{char } k \neq 2$, then $\mathcal{E}(x, k)$ is its only direct summand of this form. Moreover, it occurs with multiplicity 1.*

REFERENCES

- [AB89] A. Arabia, Cohomologie T-équivariante de la variété de drapeaux d'un groupe de Kač-Moody, *Bull. Soc. math. France*, **117** (1989), 129–165.
- [A98] A. Arabia, Classes d'Euler équivariantes et points rationnellement lisses, *Ann. Inst. Fourier*, Grenoble 48,3(1998), 861–912.
- [AB] M.F. Atiyah, R. Bott The moment map and equivariant cohomology, *Topology*, **23**(1), 1–28, 1984.
- [BL] J. Bernstein and V. Lunts. Equivariant sheaves and functors, *Lecture Notes in Mathematics*, v. 1578, Springer-Verlag, Berlin, 1994.
- [BW] T. Braden, Williamson G., Modular intersection cohomology complexes on flag varieties, *Math. Z.*, **272**(3–4)(2012), 697–727.
- [B] M. Brion, Equivariant cohomology and equivariant intersection theory, *Representation theories and algebraic geometry (Montreal, PQ, 1997)*, v. 514 of NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., pages 1–37.
- [D] A. Dimca, Sheaves in topology, *Universitext*, Springer, Berlin, 2004.
- [F] P. Fiebig, An upper bound on exceptional characteristics for Lusztig's character formula, *J. Reine Angew. Math.*, **673** (2012) 1–31.
- [G] S. Gaussent, Corrections and New Results on the Fibre of the Bott-Samelson Resolution, *Indag. Mathem.*, N.S., **14** (1), 31–33.
- [FW] P. Fiebig and G. Williamson, Parity sheaves, moment graphs and the p-smooth locus of Schubert varieties, arXiv:1008.0719.
- [H] M. Härterich, The T-Equivariant Cohomology of Bott-Samelson Varieties, arXiv:math/0412337v1.

- [I] B. Iversen, Cohomology of sheaves, *Universitext, Springer*, Berlin, Heidelberg, New York, 1986.
- [J] J. C. Jantzen, Moment graphs and representations, *Inst. Fourier, Geometric Methods in Representation Theory, Summer School Proceedings*, 2008.
- [JMW] D. Juteau, C. Mautner, and G. Williamson. Parity sheaves, *J. Amer. Math. Soc.*, **27** (4), 1169–1212 (2014).
- [Q] D. Quillen , The spectrum of an equivariant cohomology ring: I, *Ann. of Math.*, **94** (1971), 549–572.
- [S] V. Shchigolev, On decomposition of Bott-Samelson sheaves, *J. of Algebra*, **406** (2014), 376–418.

E-mail address: shchigolev_vladimir@yahoo.com